



International Research Journal of Plant Science (ISSN: 2141-5447)  
Vol. 14(3) pp. 01-06, July, 2023  
DOI: <http://dx.doi.org/10.14303/irjps.2023.17>  
Available online @ <https://www.interestjournals.org/plant-science.html>  
Copyright ©2023 International Research Journals

**Research Article**

# Allelopathic Activity of Weed *Ageratum conyzoides* on Seed Germination of Mung (*Vigna radiata*)

Versha Upadhy<sup>\*</sup>

Department of Botany, Maya Group of Colleges, Selaqui, Dehradun, India

E-mail: [dr.vershau@gmail.com](mailto:dr.vershau@gmail.com)

## Abstract

Allelopathy is the inhibition or stimulation by chemicals which is secreted by plants and microorganisms on the growth of seed germination of other plants. Use of chemical herbicide in exes amount to control the growth of weed. These herbicide increase weed resistant and create harsh environmental pollution. *Ageratum conyzoides* is one of the weed which effect the seed germination. In this experiment observe the seed germination percentage on various concentration And it was examine on different parameter like GP (Germination Percentage), GI (Germination Index), GRI (Germination Rate Index), MGT (Mean of Germination Time), GE (Germination Energy) and sum #germination. Seed germination were reduced by 90%, 60% and 40% and there is no germination on 100% concentration compared to the control, respectively.

**Keywords:** Allelopathy, Inhibition, Microorganisms herbicide, *Ageratum conyzoides* germination

## INTRODUCTION

Allelopathy is the phenomenon of inhibition or stimulation by chemicals which is secreted by one plant/microorganism on the growth on seed germination of other plants. Use of chemical herbicide in exes amount to control the growth of weed. These herbicide increase weed resistant and create harsh environmental pollution Maria. Other weed management activities are eco-friendly but cost

effective and time demanding issue throughout the world. Phytotoxic plants might help in resolving the problems created by synthetic herbicide as they possess growth retarding substances. Recently increasing interest into the researchers for the work on medicinal plants because it is easily screened from medicinal plants Kulasekaran. The increasing interest on medicinal plants could be due to either.

---

**Received:** 28-June-2022, Manuscript No. IRJPS-23-67945; **Editor assigned:** 01-July-2022, PreQC No. IRJPS-23-67945 (PQ); **Reviewed:** 15-July-2022, QC No. IRJPS-23-67945; **Revised:** 14-June-2023, Manuscript No. IRJPS-23-67945 (R); **Published:** 12-Jul-2023

---

**Citation:** Upadhy V (2023). Allelopathic Activity of weed *Ageratum conyzoides* on Seed Germination of Mung (*Vigna radiata*). IRJPS. 14: 17.

The easier screening process of phytotoxic plants from medicinal plants or the possibility to have more bioactive compounds in medicinal plants than other plants wasim. These phytotoxic plants could be used in several ways to control weeds, for example:

- Sowing/transplanting them as relay or cover crops with main crops.
- Direct application of their crude extracts as bio herbicides.
- Isolation and characterization of their active substances and using them as a tool for new natural and biodegradable herbicides development Mominul and Noguchi.

#### Nutritional value

Mung beans contain highly nutritive value. It contain 65% carbohydrate in 650 g/kg dry weight and it is rich in protein, vitamins and minerals. Around 20%-50% protein of total dry weight, with globulin 62% and albumin 27% are strongest proteins in mung beans. Mung beans are considered to be strongest protein and substantive source of dietary proteins. The proteolytic cleavage of these protein are even higher during sprouting. Bean's carbs are easy to digest ,which causes less flatulence in human compared to other forms of legumes. Both seed and sprout of mung bean produce lower calories compares to other cereals, which is more useful for obese and diabetics persons Piotr (Hoffmaster et al., 2003 and Lamsal et al., 2019).

## MATERIALS AND METHODS

#### Plant materials

*Ageratum conyzoides* is weed plant and presently a major problem for environmentalists, ecologists, farmers and animal scientists. A number of studies have been allied on its control as a weed. Its biological properties of the extracts and the constituents might provide incentive for proper evaluation of the use of the plant in medicine and in agriculture. Test conducted in mice and rats for anti-inflammatory, analgesic, antipyretic, antibacterial, anti-fungal, anti-ulcer, radio-protective activities have shown significant results without adverse side effects Riddiford (Elegami et al., 2002). Similarly, clinical trials with arthritis patients conducted with the aqueous extract of the whole plant did not show any side effect. The major constituents of the essential oil have been shown to produce precocious metamorphosis in insect larvae as well as sterility, moribund and dwarfness in adult insects (Durodola, 1977). The  $\beta$  flavonoids possess wide range of biological activities like its effects on CVS, diuresis, antiviral, spasmolytic, ant-inflammatory properties of the  $\beta$  flavonoids isolated from the plant

need to be studied. Precocenes and coumarins have been seen as fourth-generation insecticides, also need to be studied. Traditional communities in India use this species as an antidysenteric and antilithic and in Asia, South America and Africa aqueous extract of this plant is used as an antibacterial agent (Ellis et al., 1981).

*Ageratum conyzoides* L, is an annual herb with a long history of traditional medicinal uses in many countries in the world, especially in the tropical and subtropical regions. A wide range of chemical compounds including alkaloids, flavonoids, chromenes, benzofurans and terpenoids have been isolated from this species (Farooq et al., 2011). Extracts and metabolites from this plant have been found to possess pharmacological and insecticidal activities. The whole plant is only used for medicinal purposes and has a long history in the folk medicine of different countries (Islam et al., 2012). Various extracts of the plant, including water and methanol have been shown to inhibit the growth of *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, *Pseudomonas aeruginosa* and *H. pylori*. Durodola demonstrated the effectiveness of crude extract of this plant in inhibiting the growth of *Staphylococcus aureus* major wound pathogen in *in-vitro* cultures of the organism. Plants of *Ageratum conyzoides* was collected from Dehradun valley. The plant was identified by botanical survey of India, Dehradun. The roots of *A. conyzoides* were chopped off leaving the aerial part which was shredded into tiny bits (Amat et al., 2018).

#### Collection of aerial parts of *Ageratum conyzoides* material

Plant material, methanol, Whatmann filter paper, grinder, beakers, funnel, laminar air flow, autoclave, distilled water (Figure 1) (Manzano-Agugliaro et al., 2013).



Figure 1. *Ageratum conyzoides*.


#### Extraction procedure method

The plant material was washed gently with tin twenty to remove any dirt and to free it from any microbe and was air-dried under shade for a week. The sample was powdered with an electric grinder into a coarse form and stored in airtight containers (Table 1) (Pell et al., 1998 and Aktar et al., 2009).

**Table 1.** The plant material containers

Plant material	Fresh weight	Dried weight
<i>Ageratum conyzoides L</i>	500 grams	150 grams

After grinding, 30 gm of plant material was extracted in 130 ml of methanol for 24 hours. The extracts were filtered through Whatmman filter paper and were evaporated to dryness using a hot plate at a much reduced temperature (40°C). The residues obtained were dissolved in methanol. The weights of the extract was determined and stored below ambient temperature (Figure 2) (Ndip et al., 2007).



**Figure 2.** (a) shredding of plant parts, (b) and (c) preparation of crude methanol extract (d) methanolic extract of plant (green in colour).

### Germination bioassay

Plant extract was diluted into small amount of distil water to prepare four assay concentration 25% 50% 75% and 100% and then was spray on filter paper No 1 in petri dishes. Ten seeds of mung was use for germination (Antonisamy et al., 2017).

Ten seeds of mung placed on the filter paper in Petri dishes. Control Petri dishes were also maintained in each experiment using 10, that is, without plant extracts. The Petri dishes were then incubated in dark at 25°C. Germination was measured at every 0.5-day interval up to 7 days (the time when no further seeds germinated) and was considered when the radical emerge by rupturing the seed coat as per (Figure 3).

Eight germination indices, that is, Germination Percentage (GP), Germination Index (GI), Germination Energy (GE), mean Germination Time (MGT), were calculated GP index indicated the total germination percent of a seed lot after certain period of time when germination became constant. As it is measured by total germination relative to total number of seeds set for germination, GP cannot explain the delayed germination. In contrast, GI is a measure of both

percentage and speed of germination and assigns maximum arithmetic weight to seeds that germinate during first count and less weight to those that germinate later. The higher the GI, GE, values compared to control, the lower the inhibition and vice versa. But the meaning is reversed for MGT. Germination Percentage (GP)=No of seed germinated/Total no of seed tested\*100 he Germination Index (GI) was calculated as described in the association of official seed analysts by following formula:

$$\text{Germination Index} = \sum (GT / Tt) \text{ or } \left[ \frac{\text{No. of germinated seed}}{\text{Days of first count}} \right] + \dots + \left[ \frac{\text{No. of germinated seed}}{\text{Days of final or last count}} \right]$$

Germination Energy (GE)=No of seed germinated on day/Total no of seed tested\*100

Germination Rate Index (GRI)=(G1/1)+(G2/2)+(G3/3)+(Gi/i)+....

Mean Germination Time (MGT)=MGT= $\sum(n \times d)/N$

n=number of seeds germinated on each day, d= number of days from the beginning of the test and N= total number of seeds germinated at the termination of the experiment.



**Figure 3.** Seeds germinated at the termination.

### Growth bioassay

The Petri dishes and the extracts were prepared as described above. Ten pregerminated seeds of mung, (germinated in the darkness at 25°C for 1-3 days after overnight soaking) were placed on the filter paper in Petri dishes. The shoot and root lengths of each seedling were measured after incubation in dark condition for 2 days at 25°C. Control Petri dishes were also maintained as germination bioassay (Gulewicz et al., 2008).

### Statistical analysis

The bioassay experiments were conducted as Completely Randomized Design (CRD) with three replications. The experiments were repeated twice to

avoid any experimental error. The data calculated on different parameter for its verification.

## RESULTS AND DISCUSSION

*Ageratum conyzoides* aqueous extract was used in an allelopathic study to examine the potential allelopathic effects of its flower, root, stem, and leaf extracts on the growth of Mung bean seedlings, percentage germination, and seedling evaluation at treatment concentrations of 10 mg/ml, 20 mg/ml, 50 mg/ml, and 100 mg/ml, respectively. After 10 days in vitro, the effects of concentration on seed germination, root length and seedling height of mung bean were assessed. Results obtained demonstrated substantial differences in all three parameters investigated between the various concentrations of aqueous leaf extracts employed when compared with the control.

*Ageratum conyzoides* aqueous leaf extract showed a significant reduction in seed germination, root growth, and seedling height in all treatments, though at varying degrees ranging from mild to severe effects. In contrast, aqueous Mung bean extract showed an increase in seed germination, root length, and plant height. *Ageratum conyzoides* aqueous extracts

decreased the length of the mung radicle. The length of the mung radicle, biochemical parameters and cumulative germination percentage were all affected negatively by the weed plant extract's growth-inhibiting properties. According to the current study, *Ageratum conyzoides* plants have a high potential for allelopathy against *Vigna radiata*.

Allelopathy is the creation of one or more biochemicals by an organism that influence the germination, growth, survival, and reproduction of other organisms is a biological phenomenon. The target organisms and the community may be impacted favourably (positive allelopathy) or negatively (negative allelopathy) by these allelochemicals, which are also known as biochemical (Sullivan et al., 2000). Allelochemicals are a subset of secondary metabolites that are not necessary for the allelopathic organism's metabolism (*i.e.*, growth, development, and reproduction). An essential component of plant defence against herbivory is the use of allelochemicals having adverse allelopathic effects. Allelochemical production is affected by both biotic and abiotic variables, including temperature and pH (Table 2 and Figure 4) (Del Rosario and Flores, 1981).

**Table 2.** Calculation on seed germination 25% concentration.

Seed Mung	Day 1	Day2	Day3	Day4	Day5	Day6	Day7	Total no of seed	Sum # germ	GRI	MGT	GP	GI	GE
<b>25% concentration</b>														
Trial on 10 seed	10	10	10	10	10	10	10	10						
R1	0	0	3	4	6	7	9	10	29	15.99	17.77	90	5.65	290
R2	0	0	4	5	7	7	8	10	31	16.57	20.62	80	4.89	310
R3	0	0	3	3	4	8	9	10	27	16.74	16.88	90	5.16	270
<b>50% concentration</b>														
Trial on 10 Seed	10	10	10	10	10	10	10	10						
R1	0	0	3	3	4	5	6	10	21	11.95	18	60	4.24	210
R2	0	0	2	5	5	5	5	10	22	9.64	23.2	50	4.46	220
R3	0	0	4	4	4	6	6	10	24	12.7	21	60	4.99	240
<b>75% concentration</b>														
Trial on 10 seed	10	10	10	10	10	10	10	10						
R1	0	0	2	4	4	4	4	10	18	8.84	23.5	40	3.7	180
R2	0	0	3	3	3	4	4	10	17	8.73	22	40	3.58	170
R3	0	0	3	3	4	4	4	10	18	8.93	23.25	40	3.78	180



100% concentration														
Trial on 10 seed	10	10	10	10	10	10	10	10						
R1	0	0	0	0	0	0	0	10	0	0	0	0	0	0
R2	0	0	0	0	0	0	0	10	0	0	0	0	0	0
R3	0	0	0	0	0	0	0	10	0	0	0	0	0	0
Controlled														
Trial on 10 seed	10	10	10	10	10	10	10	10						
R1	1	2	4	5	7	8	10	10	37	21.6	19	100	8.74	370
R2	2	3	5	6	7	7	10	10	40	23.51	19.4	100	10.7	400
R3	1	3	4	6	6	8	9	10	37	20.72	20.37	100	9.15	370




Figure 4. The chemical components analysis of different extracts.

## CONCLUSION

The chemical components analysis of different extracts of *Ageratum conyzoides* In our study, I found that *Ageratum conyzoides* whole plant significantly inhibited the germination and reduced the varieties and biomass of weeds in the field, when it was applied as a fertilizer originally. Therefore, I think that certain allelochemicals present in might inhibit the growth of weeds. To investigated the possible allelochemicals in *Ageratum conyzoides*, solvents (methanol) was used to extract the metabolites in *Ageratum conyzoides* leaves. The crud extract was analysed as shown in Table 1 that extract reduced germination of weed species tested. Seed Germination were reduced by 90%, 60% and 40% and there is no germination on 100% concentration compared to the control, respectively.

## REFERENCES

Hoffmaster AL, Fujimura K, McDonald MB, Bennett MA (2003). An automated system for vigor testing three-day-old soybean seedlings. *Seed Sci Technol.* 31: 701-713.

Lamsal A, Devkota MP, Shrestha DS, Joshi S, Shrestha A (2019). Seed germination ecology of *Ageratum houstonianum*: A major invasive weed in Nepal. *PLoS One.* 14: 0225430.

Elegami AA, El-Nima EI, Tohami ME, Muddathir AK (2002). Antimicrobial activity of some species of the family *Combretaceae*. *Phytother Res.* 16: 555-561.

Durodola JI (1977). Antibacterial property of crude extracts from a herbal wound healing remedy- *Ageratum conyzoides* L. *Planta Med.* 32: 388-390.

Ellis RH, Hong TD, Roberts EH (1981). The influence of desiccation on cassava seed germination and longevity. *Ann Botany.* 47: 173-175.

Farooq M, Jabran K, Cheema ZA, Wahid A, Siddique, KH (2011). The role of allelopathy in agricultural pest management. *Pest Manag Sci.* 67: 493-506.

Islam AKMM, Kato-Noguchi H (2012). Allelopathic potentiality of medicinal plant *Leucas aspera*. *Int J Sustain Agric.* 4: 01-07.

Amat I, Desouhant E, Gomes E, Moreau J, Monceau K (2018). Insect personality: What can we learn from metamorphosis? *Curr Opin Insect Sci.* 27: 46-51.

Manzano-Agugliaro F, Alcayde A, Montoya FG, Zapata-Sierra A, Gil C (2013). Scientific production of renewable energies worldwide: An overview. *Renew Sust Energ Rev.* 18: 134-143.

Islam AM, Kato-Noguchi H (2013). Plant growth inhibitory activity of medicinal plant *Hyptis suaveolens*: Could allelopathy be a cause? *Emir J Food Agric.* 692-701. [Crossref] [Google Scholar]

Pell, M., Stenberg, B., Torstensson, L. (1998). Potential denitrification and nitrification tests for evaluation of pesticide effects in soil. *Ambio.* 24-28.

Aktar MW, Sengupta D, Chowdhury A (2009). Impact of pesticides use in agriculture: Their benefits and hazards. *Interdiscip Toxicol.* 2: 1-2.

Ndip RN, Tarkang AEM, Mbullah SM, Luma HN, Malongue A, et al (2007). *In vitro* anti-*Helicobacter pylori* activity of extracts of selected medicinal plants from North West Cameroon. *J Ethnopharmacol.* 114: 452-457.

## 6 Int. Res. J. Plant. Sci.

- Antonisamy, P., Dhanasekaran, M., Kim, HR., Jo, SG., Agastian, P., et al (2017). Anti-inflammatory and analgesic activity of ononitol monohydrate isolated from *Cassia tora* L. in animal models. Saudi J Biol Sci. 24: 1933-1938. [Crossref] [Google Scholar] [PubMed]
- Gulewicz, P., Martinez-Villaluenga, C., Frias, J., Ciesiolka, D., Gulewicz, K., et al (2008). Effect of germination on the protein fraction composition of different lupin seeds. Food Chem. 107: 830-844.
- Sullivan JP, Jassim O, Fahrbach SE, Robinson GE (2000). Juvenile hormone paces behavioral development in the adult worker honey bee. Horm Behav. 37: 1-14.
- Del Rosario RR, Flores DM (1981). Functional properties of four types of mung bean flour. J Sci Food Agric. 32: 175-180.