



*Full Length Research Paper*

# **An assessment of the nutritional knowledge, practice and status of adult HIV/AIDS patients attending an art centre in Jos, north central Nigeria**

**\*Banwat, M.E., Yakubu N.W., Olalude E.O., Ogunsakin, J.A.**

Department of Community Health University of Jos, Plateau State Nigeria.

Department of Community Health, Faculty of Medical Sciences University of Jos, Nigeria.

\*Corresponding authors email: [mathildabanwat@yahoo.com](mailto:mathildabanwat@yahoo.com); Tel: 234-8036133983

## **ABSTRACT**

The intimately multidirectional relationship between nutrition and HIV/AIDS has been established. Although adequate nutrition can neither cure nor prevent the infection, it plays a major role in the maintenance and improvement of the immunological status of PLWHA thus delaying disease progression and improving patients' quality of life. This study aimed at assessing the factors affecting knowledge and practice of adequate nutrition among adult HIV/AIDS patients attending an AIDS Out-Patient clinic in Jos, North-Central Nigeria. This was a cross-sectional descriptive study where 250 patients on treatment in APIN Clinic were selected using systematic sampling technique. Structured interviewer-administered questionnaires were used to gather data from them which were collated and analyzed using EPI info version 3.5.3 Statistical software. Chi-square statistical test was used to assess association and a p-value of less than or equal to 0.05 was considered to be statistically significant. Majority of the respondents (55.9%) felt protein was the most important food nutrient in their diet. Many of the respondents (48%) reported spending up to 25-50% of their monthly income to ensure they have adequate nutrition. A small proportion of the respondents (22.9%) reported early satiety as the main factors affecting their daily dietary intake while the cost of a food item was the major hindrance to ingesting a preferred food group. Forty five percent of the respondents reported fruits and vegetables as the major components of their daily diet. There was a statistically significant relationship between the BMI of the respondents and the duration on ART prior to the study. The knowledge and practice of adequate nutritional intake was fairly good among studied HIV/AIDS patients. It however needs to be improved through more involvement of health workers, the media, as well as governmental and non-governmental organizations in nutrition education and poverty alleviation.

**Keywords:** Nutritional status, HIV/AIDS, Nutritional Practices.

## **INTRODUCTION**

Human Immune-deficiency Virus (HIV) infection and Acquired Immune Deficiency Syndrome (AIDS) are caused by the Human Immune-deficiency Virus which has various serotypes. Since the isolation of the first confirmed case of the disease in 1981, HIV/AIDS has become a global health problem, with the bulk of the disease borne by Sub-Saharan Africa. (Report on the global HIV/AIDS pandemic 2009). Many HIV-related conditions affect and are affected by the body's nutritional status (Bartlett, 2003).

HIV and nutrition are interrelated and as antiretroviral

drug become increasingly available in the poorest parts of the world, critical questions are emerging about how well the drugs work in people if they are short of food, and for those already receiving treatment, side effects such as body fat changes are a daily concern. (Bartlett, 2003 and Association of Nutrition Services Agencies (ANSA)) Adequate nutrition refers to intake of a diet which meets the specific nutritional needs for the specific individual for that specific period in time. (Association of Nutrition Services Agencies (ANSA)). The sole aim of adequate nutrition is to meet the growth and

developmental demands of the unique, specific individuals' body. (Walsh et al., 2003) Inadequate nutrition in people with HIV infection may result from many factors including nausea, vomiting and anorexia that may prevent adequate intake of nutrients and medications; diarrheal infections that prevent absorption of nutrients and medications; Oral health conditions that interfere with chewing or tasting food like Oral Candidiasis in patients who present late; systemic illness (including HIV itself) that create a catabolic state; and psychological conditions such as depression that impair patients' ability to nourish themselves. (Steinhart, 2011 and Association of Nutrition Services Agencies (ANSA)) In addition, financial constraints may limit patients' access to nutritious food (Walsh et al., 2003).

Adequate nutrition helps to maintain and improve the nutritional and immunological status of a person with HIV/AIDS and delay the progression from HIV to AIDS-related diseases (Steinhart, 2011). It can therefore improve the quality of life of PLWHA. Adequate nutrition will complement the effects of antiretroviral therapies and will help to maintain body weight and fitness, as well as improve the performance of the immune system already compromised by the infection (Steinhart, 2011). Whereas starving people tend to lose fat first, the weight lost during HIV infection tends to be in the form of lean tissue such as muscle mass. This means there may be changes in the makeup of the body even if the overall weight stays the same (Evidence lessons and recommendations for action, April 2005 issue and World Health Organization (WHO), African Regional report 2008) One factor behind HIV-related weight loss is increased energy expenditure (European Collaborative Study 2003 and Batterham, 2005) Many studies indicate that people with HIV tend to burn around 10% more calories while resting, compared to those who are uninfected (Evidence lessons and recommendations for action, April 2005 issue and Batterham, 2005). Nutritional control of weight can be effectively used to balance out these body changes in infected patients.

Current antiretroviral drug treatments control HIV infection and prevent severe wasting. In addition, some antiretroviral drugs have been linked to lipo-dystrophy, as well as lipid abnormalities by raising Low Density Lipoprotein (LDL) cholesterol, lowering High Density Lipoprotein (HDL) cholesterol and raising triglyceride levels in the blood (Batterham, 2005 and Sabin et al 2008). One study in Malawi found that patients with mild malnutrition (a Body Mass Index (BMI) of 17.00 - 18.49kg/m<sup>2</sup>) were twice likely to die in the first three months of treatment; and for those with severe malnutrition (BMI less than 16.00kg/m<sup>2</sup>), the risk was six times greater than those of healthy body weight (BMI ranging between 18.5 - 24.99kg/m<sup>2</sup>) (Paton, 2003).

Since the beginning of the pandemic, over 30 million people have died from AIDS-related causes globally (Report on the Global HIV/AIDS pandemic 2

010). In Nigeria, an estimated 4.6% of the populations are living with HIV/AIDS (The United Nations General Assembly Special Session Country Progress report: Nigeria 2010) Although HIV prevalence is much lower in Nigeria than in other African countries the enormous size of Nigeria's population makes this prevalence to be a large number. The HIV pandemic has also significantly compromised the food security of affected households and communities, reducing the availability of productive labor, diverting income, depleting savings and productive assets and impeding intergenerational knowledge transfer (Human Development Reports (2007/2008)).

This study aimed at assessing the factors affecting knowledge and practice of adequate nutrition among adult HIV/AIDS patients attending an AIDS Out-Patient clinic in Jos, North-Central Nigeria.

## METHODS AND MATERIALS

### Study Area

The study was conducted in the AIDS Prevention Initiative in Nigeria (APIN) centre within Jos University Teaching Hospital (JUTH) in Jos, Plateau State, North-Central Nigeria. APIN is a Public-Private Partnership funded organization, which is involved in the provision of HIV/AIDS prevention, treatment and care services. Other services it renders include: Prevention of mother to child transmission (PMTCT) of HIV, Paediatric HIV care and treatment, Tuberculosis/HIV care and treatment, Laboratory infrastructure procurement and supply and Training and human resource capacity building.

APIN centre is located in Jos North Local Government Area (LGA) of Plateau State, North-Central Nigeria, within the temporary site of the Jos University Teaching Hospital (JUTH).

### Study Design

This was a cross-sectional, descriptive study, to assess the knowledge and practice of adequate nutrition among HIV/AIDS patients in APIN Centre, Jos.

### Study Population

The majority of patients at APIN centre, come from within Plateau state, with a significant number also coming from neighboring states within the North-Central part of Nigeria including Benue, Kaduna and Nassarawa states. The bulk of patients at APIN centre are adult HIV patients although a significant number of children also attend the paediatric clinics. The social status of patients at APIN centre cut across all social strata with most of the adults being literate and employed in professional, skilled and

manual labor; A Professional occupation has an approved standard curriculum of training for students in institutions (Lawyer, Medical doctor, Nurse), a skilled profession needs a period of skill acquisition as an apprentice under an instructor but has no curriculum for training (Tailor, Welder, Motor mechanic) while a manual laborer needs no specific period under an instructor to be in the trade (a Night watchman, farmer, petty trader).

### Inclusion/Exclusion criteria

The target population of the study comprised of adult patients (18years and above) attending APIN centre for at least three months prior to the study; all subjects were either on ART or had been on ART within the past 12months of the study. Patients less than 18 years of age and over 65 years, indigent patient, as well as those registered in the APIN clinic, less than three months prior to the study were excluded from the study. The elderly (aged above 65years) excluded from the study since aging on its own is a predisposing factor for malnutrition and could act as a confounder. The use of BMI to assess nutritional status is also unreliable due to skeletal deformities with age which affect height.

### Sample Size Determination

A minimum sample size was determined using the formula:

$$N = \frac{Z^2 pq}{d^2}$$

Where:

N = Minimum sample size

Z = Standard deviation at 95% confidence interval = 1.96

p = Anticipated proportion/population of PLWHA with appropriate knowledge and practice of adequate nutrition = 20.9% (0.209) from a similar survey (Tang, 1993).

q = Complementary probability =  $1 - p$   
 $= 1 - 0.209$   
 $= 0.791$

d = Error margin/tolerance = 5% = 0.05

Hence:  $N = \frac{(1.96)^2 \times 0.209 \times 0.791}{(0.05)^2}$

$N = \frac{0.6351}{0.0025} = 244.04$

N ~ 250

The minimum sample size was therefore 250 patients.

### Sampling Technique

The sampling method used was systematic sampling; of the equal – probability modality. This sampling interval was elucidated using the formula:

$$K = \frac{N}{n}$$

Where:

K = sampling interval by which every  $K^{\text{th}}$  element/subject was selected from the sampling frame.

N = population size of patients =  $1000 \times 2 = 2000$  (in four weeks)

n = sample size = 250 patients

Hence  $K = \frac{2000}{250} = 8$

Therefore, within 4 weeks of sampling, every 8<sup>th</sup> patient was selected from the entire sampling frame of 2000 patients to comprise the 250 patients as sample size, given the logical homogeneous composition of the population.

### Instruments for Data Collection

Data was collected by interview-administered structured questionnaires which was framed by the researcher from the objective of the research and was up of five sections: Section A: assessed the Socio-demographic data of the patients

Section B: Assessment of knowledge of adequate nutrition

Section C: Assessment of practice of adequate nutrition

Section D: Assessment of the determinants and deterrents to adequate nutrition.

Section E: Others – including nutritional assessment by anthropometric measurements.

Weighting scales and meter rules were used to obtain weights (in kilograms) and heights (in meters) of subjects respectively, using standard procedure: Subjects were weighed twice with minimal clothing and the average of the reading recorded while they stood against a wall with their heel, buttocks and vortex against the wall for their heights to be measured in meters. Their BMI was then calculated (Weight in kilograms/Height in meters<sup>2</sup>) and recorded.

### Data Analysis

Data collected was collated and analyzed using EPI info version 3.5.3 Statistical software. Significant relationship between BMI of respondents and the duration of HAART was evaluated using Chi-squared ( $\chi^2$ ) test. The p value of  $\leq 0.05$  was taken as statistically significant.

### Ethical Considerations

Ethical approval for the study was given by the Research and Ethics Committee of the Jos University Teaching

**Table 1.** SOCIO-DEMOGRAPHICS OF THE RESPONDENTS:

The mean age of respondents was  $37.8 \pm 8.9$  years with the median duration on ART being 4.38 years. The highest proportion of respondents were aged between 30 and 39 years (44.4%), female (52.8%), skilled workers (31.2%) and married 54.8%

VARIABLE	FREQUENCY (n = 250)	PERCENTAGE
<b>AGE(YEARS)</b>		
18-29	39	15.6
30-39	111	44.4
40-49	74	29.6
50-59	26	10.4
<b>SEX</b>		
Males	118	47.2
Females	132	52.8
<b>HIGHEST EDUCATIONAL STATUS ATTAINED</b>		
None	5	2.0
Primary	23	9.2
Secondary	90	36.0
Tertiary	132	52.8
<b>OCCUPATION</b>		
Students	40	16.0
Professional	61	24.4
Skilled worker	78	31.2
Unskilled worker	36	14.4
Unemployed	35	14.0
<b>MARITAL STATUS</b>		
Single	66	26.4
Married	137	54.8
Divorced	10	4.0
Separated	3	1.2
Widow/Widower	34	13.6

**Table 2.** PERCEPTION OF THE MOST VITAL FOOD NUTRIENT IN THEIR DIET

Majority of the respondents (55.9%) felt protein was the most important food nutrient in their diet while only 2.9% felt water was the most important. Majority of the respondents (54.5%) cited building the immune system as the reason for their consideration of a food nutrient as important while only 6.8% felt it helped them look good

Variable	Frequency (n=250)	Percentage
<b>Most Important Food Nutrient</b>		
Protein	117	46.8
Minerals/Vitamins	66	26.4
Carbohydrates	13	5.2
Fats/Oils	7	2.8
Water	6	2.4
Don't Know	41	16.4
<b>Reason for choice of Nutrient</b>		
It builds immunity	114	45.6
It improves wellbeing	29	11.6
It provides energy	25	10.0
It helps drugs work well	21	8.4
It enables weight gain	14	5.6
Others*	47	18.8

Others\* = Cleanses the body, repairs damages made by disease, no reason given.

Hospital (JUTH). A letter of approval was also obtained from APIN centre to conduct the study. A letter of identification from the Department of Community Health was given to the research team. Prior to administration of

the questionnaires, written informed consent was obtained from each patient after being assured of the the participants included free nutritional counseling and

**Table 3.** PROPORTION OF MONTHLY INCOME SPENT ON NUTRITION

Many of the respondents (48%) reported spending up to 25-50% of their monthly income to ensure they have adequate nutrition

Proportion of Monthly income (%)	Frequency (n=250)	Percentage
<25	41	16.4
25-50	120	48.0
51-75	68	27.2
>75	21	9.4
TOTAL	250	100

**Table 4.** FACTORS AFFECTING DIETARY INTAKE

About 23% of the respondents reported early satiety as the main factors affecting their daily dietary intake while the cost of a food item was the major hindrance to ingesting a food item even if a respondent wanted to eat it. Forty five percent of the respondents reported fruits and vegetables as the major components of their daily diet. Only 1.6% reported eating a lot of fatty and oily foods

Variable	Frequency (n=250)	Percentage
<b>Factors affecting quantity of meals</b>		
Early satiety	48	22.9
Depression/stress	45	21.4
Adverse effect of drugs	25	11.9
Diarrhea	18	8.6
Constipation	17	8.1
Difficulty coordinating meals with drugs	15	7.1
Heart burn	12	5.7
Nausea/vomiting	10	4.7
Pain on swallowing	7	3.3
Use of supplements	7	3.3
Difficulty with swallowing	6	2.9
<b>Factors Quality of meals</b>		
Availability of food items	45	18.0
Geographical access	53	21.2
Cost of food items	99	39.6
Seasonality of food items	49	19.6
Social issues	41	16.4
None given	38	15.2
<b>Major component of daily diet</b>		
Cereals	68	27.2
Meat/Fish/Milk/Legumes	65	26.0
Fruits /Vegetables	114	45.6
Fats/Oils	3	1.2

confidentiality of volunteered information. Incentives to education on the research topic.

### Limitations to the Study

1. The nature of the study, being a "snapshot" of the nutritional characteristics of the patients, hindered any prospective/follow-up activities with appropriate documentation.
2. Recall bias remains a limitation since the subjects had to remember components of their nutritional intake as well as other discrete pieces of information

## RESULTS AND DISCUSSION

A good knowledge of nutritional requirements for an individual with a chronic disease has been found to be vital for proper management of the disease as well as prevention of complications (Evidence, lessons and recommendation for action (2006): Tanzania AIDS Management Training Programme 2011 and Erica, 2012). Majority of the respondents (55.9%) in this study considered protein to be the most important food nutrient they need to ensure an adequate diet and to ensure they stay strong despite their diseased state. This suggests that a good proportion of them have an understanding of the fact that HIV/AIDS is an immune deficiency disease

**Table 5.** RELATIONSHIP BETWEEN BODY MASS INDEX (BMI) AND DURATION OF USING ANTIRETROVIRAL THERAPY (ART). There was a statistically significant relationship between the BMI of the respondents and the duration since commencement of ART prior to the study.

BMI	Duration of HAART (Years)				Total Freq (%)
	< 1 Freq (%)	1 – 5 Freq (%)	6 – 10 Freq (%)	>10 Freq (%)	
<18.5 (Underweight)	2 (3.6)	1(1.1)	3 (4.4)	1 (3.1)	7 (2.8)
18.5-24.9 (Normal weight)	31 (55.4)	60(63.8)	26 (38.2)	15 (46.8)	132 (52.8)
25.0-29.9 (Overweight)	15 (26.8)	20 (21.3)	29 (42.6)	11 (34.4)	75(30.0)
30.0-39.9 (Obesity)	7 (12.5)	12 (12.7)	10 (14.8)	5 (15.7)	34(13.6)
>40.0 (Morbid obesity)	1 (1.7)	1(1.1)	0	0	2(0.8)
TOTAL	56	94	68	32	250

$$\chi^2 = 15.18; df = 12; P = 0.0231$$

and the role of protein in the production and functioning of the immune system. This perception was further strengthened by majority of them giving the principal reasons for placing priority on that particular food nutrients included its function in building and strengthening of the immune system and improvement of health. A study in Anambra state of Nigeria, among pregnant women also showed an above-average knowledge score regarding food classes required in pregnancy and the usefulness of protein in ensuring a healthy mother and baby at the end of pregnancy (Ihekoronye 2011). Nutrition guidelines for HIV/AIDS patients explains the importance of protein in building muscles, organs and a strong immune system (Evidence, lessons and recommendation for action (2006) and Erica, 2012). Furthermore nutritional counseling given to HIV/AIDS patients in clinics (like the study area) provides the needed information on the importance of food nutrients.

In this study, most of the participants reported an increase in cost and geographical inaccessibility of food items as the major constraints to adequate dietary intake in terms of quantity and quality of meals ingested. This could be associated to the crises in Jos which make food cultivation and transportation difficult and expensive. This is in consonance with a study by Nelson et al in Delta State, Nigeria where inadequate funds and location of markets were major determinants of food items ingested by the studied population (Nelson, 2012). It is however slightly different from findings among studied pregnant women in Imo state of Nigeria, where cultural factors like food taboos were a major constraint to ingestion of certain food items; as well as cost (Ihekoronye, 2011). The uniformity of cost as a constraint in these and many other studies may be attributed to the general dire economic and financial constraints experienced nationally and globally with its attendant inflation of food prices. It can also be explain by the fact that the respondents in this study who are majorly workers have a restricted schedule and apportioned time for eating with limited choices of food varieties during working hours.

In this study, a large proportion of respondents spend between 25-50% of their monthly income on nutrition. This finding is similar to that calculated by the United States Development Agency for Nigeria of 40.1%.<sup>21</sup> It is also comparable to that of other developing countries; Cameroon (38.5%), India (35.8%), China (33.9%) but far higher than findings in developed countries like United Kingdom (8.9%), United States of America (6.8%) and Canada (9.1%) (Maia-Leite, 2012). This difference between developing and developed economies can be explained by the poverty in developing countries which means the meager income in most homes is used to meet basic physiological needs. It can also result from the fact that an average Nigerian family size is larger than what is found in developed countries; this increases the proportion of income required to meet nutritional needs (Adebayo, 2012).

Approximately 53% of the respondents (using Body Mass Index, BMI), were of normal weight, 30% overweight and 14.4% obese; with only 2.8% being underweight. There was a statistically significant relationship noted between BMI and duration of commencement of antiretroviral therapy (ART). This association may be explained by the ability of the drugs to decrease resting-energy expenditure in the respondents which is reportedly about 10% and 20% higher in asymptomatic and symptomatic HIV patients respectively (Maia-Leite, 2010 And Tapam et al 2008). The drugs help to decrease the basal metabolic rate commonly responsible for weight loss, thus contributing to replenishment of muscle bulk that translates to attaining normal or even overweight. A similar relationship between progression to Obesity and duration of ART was reported among Brazilian HIV/AIDS patients (United States Development Agency/Economic Research Service 2012).

## CONCLUSION AND RECOMMENDATIONS

It was concluded from this study that knowledge of adequate nutrition was fairly good among the studied APIN patients. Most of the respondents had good



practices of adequate nutrition. The major factors found to affect dietary intake of required food items was cost of the foodstuff and accessibility (geographical and seasonal). There was also a statistically significant relationship between body mass index BMI and duration of usage of antiretroviral therapy ( $P = 0.0231$ ).

## REFERENCES

- Adebayo OO(2012). Family income among small scale farmers; a panacea for household security in Oyo state, Nigeria. *Research in Humanities and Social Sciences*.2012;2(7):32-36.
- Association of Nutrition Services Agencies (ANSA) ANSA Nutrition guidelines and fact sheets. Available at [www.aidsetc.org](http://www.aidsetc.org) Last accessed: 6<sup>th</sup> December, 2011
- Bartlett JG(2011).Introduction: Integrating nutrition therapy into medical management of Human Immunodeficiency Virus. *Clinical Notes on Infectious Diseases*, 2003; 36(12):s51. Available at [www.aidsetc.org](http://www.aidsetc.org) Last accessed: 6<sup>th</sup> December,
- Batterham MJ(2005). Investigating heterogeneity in studies of resting energy expenditure in persons with HIV/AIDS: a meta-analysis. *American Journal of Clinical Nutrition*. 2005;81(3):92-94
- Erica C(2012).Barriers to equitable access to quality health information. *Journal of Medicine* (5): pp. 117-123. Available at <http://e-health.org/articles.html> Last accessed: 21<sup>st</sup> May, 2012.
- European Collaborative Study, Height, weight and growth in children born to mothers with HIV-1 infection in Europe. *Pediatrics*,2003;3(1):81-83
- Ihekoronye CN, Ibekano CU(2011). Nutritional Knowledge and Beliefs of Pregnant women in Aboh-Mbaise LGA of Imo State. *Nigerian Journal of Nutritional Sciences*. 2011;31(1) 36-39.
- Maia-Leite LH , Mattias-Mannho-Sampaio AB(2010). Progression to Overweight, Obesity and associated factors after ART initiation among Brazilian Persons with HIV/AIDS. *Hospital Nutrition*. 2010; 25(4): 635-640.
- Nelson E(2012) Access to health information by people living with HIV/AIDS in Nigeria (2010) Available at [www.unlib.unl.edu/LPP/edewor.htm](http://www.unlib.unl.edu/LPP/edewor.htm) Last accessed: 21<sup>st</sup> May, 2012.
- Paton NI(2006). The impact of malnutrition in survival and the CD4 count response in HIV-infection, Starting antiretroviral therapy (July 2006). *HIV Medicine*. 2006;7(5):513-515.
- Sabin,CA, Worm SW, Weber R, Reiss P, El-Sadr W(2008). Use of nucleoside reverse transcriptase inhibitors and risk of myocardial infarction in HIV – infected patients enrolled in the D. A. D. study : Multi – cohort collaboration. *The Lancet*.2008;371 (9622): 1417-1426.
- Steinhart CR(2011).HIV – associated wasting in the era of Highly Active Antiretroviral Therapy (HAART): A Practice – based approach to diagnosis and treatment. *AIDS Reader*, 2011;11(1):557-567.
- Tang AU(2006). Dietary micronutrient intake and risk of progression to Acquired immune Deficiency syndrome (AIDS) in HIV type 1 infected homosexual men *American Journal of Epidemiology*, 1993;138(11):937-941
- Tanzania AIDS Management Training Programme, *Journal of Society for women and AIDS in Dar es Salaam, Tanzania* (2011) Available at [www.ppu.go.tz](http://www.ppu.go.tz) Last accessed: 21<sup>st</sup> May, 2012
- Tapam N, Gatongi P, Rotich J, Kimaiyo S (2008).Factors affecting antiretroviral drug adherence among HIV/AIDS patients at Moi teaching and referral hospital, Eldoret Kenya. *East African Journal of Public Health*, 2008;5(2):74-78 Available at [www.ncbi.nlm.gov/pubmed/19024414.html](http://www.ncbi.nlm.gov/pubmed/19024414.html) Last accessed: 1<sup>st</sup> February, 2012.
- The Joint United Nations Programme on HIV/AIDS (UNAIDS), Report on the global HIV/AIDS pandemic (2009). Available at [www.unaids.org/en/media/unaids...GARPR\\_2013\\_guidelines\\_en.pdf](http://www.unaids.org/en/media/unaids...GARPR_2013_guidelines_en.pdf).Last accessed: 6<sup>th</sup> December, 2011
- The Joint United Nations Programme on HIV/AIDS (UNAIDS), Report on the Global HIV/AIDS pandemic (2010) Available at [www.unaids.org/documents/20101123\\_globalreport\\_em.pdf](http://www.unaids.org/documents/20101123_globalreport_em.pdf). Last accessed: 6<sup>th</sup> December, 2011
- The United Nations General Assembly Special Session Country Progress report: Nigeria (2010) Available at [www.unaids.org/.../countryprogressreports/2012countries/Nigeria](http://www.unaids.org/.../countryprogressreports/2012countries/Nigeria). Last accessed: 6<sup>th</sup> December, 2012.
- United Nations Development Programme (UNDP), Human and income poverty: developing countries, Chapter 3: Human Development Reports (2007/2008), Available at [www.undp.org/content/dam/undp/.../HDR2013%20Report%20Englis h.pdf](http://www.undp.org/content/dam/undp/.../HDR2013%20Report%20Englis h.pdf). Last accessed: 6<sup>th</sup> December, 2011
- United States Development Agency/Economic Research Service. Annual Income Spent on Food. Available at [wsm.wsm.edu/researcher/WSMaugii\\_billions.pdf](http://wsm.wsm.edu/researcher/WSMaugii_billions.pdf). Last accessed 5<sup>th</sup> February 2012.
- Walsh C, Dannhauser A, Joubert G(2011). Impact of a nutrition education programme on nutrition knowledge and dietary practices of lower socioeconomic communities in the Northern Cape. *South African Journal of Clinical Nutrition*, 2003;16(1):89-95 Available at [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) Last accessed: 6<sup>th</sup> December.
- World Health Organization (WHO) Consultation on Nutrition and HIV/AIDS in Africa: Evidence, lessons and recommendation for action (2006) Available at
- World Health Organization (WHO), African Regional report (2008), Available at [www.afro.who.int/index.php?option=com\\_docman&task=doc](http://www.afro.who.int/index.php?option=com_docman&task=doc). Last accessed: 6<sup>th</sup> December, 2011
- World Health Organization (WHO), Consultation on Nutrition and HIV/AIDS in Africa: Evidence lessons and recommendations for action, April 2005 issue. Available at [www.who.int/nutrition/topics/consultation\\_nutrition\\_and\\_hiv/htm l](http://www.who.int/nutrition/topics/consultation_nutrition_and_hiv/htm l). Last accessed: 6<sup>th</sup> December, 2011
- [www.who.int/nutrition/topics/Paper\\_1\\_Macronutrients\\_bangkok.pdf](http://www.who.int/nutrition/topics/Paper_1_Macronutrients_bangkok.pdf). Last accessed: 1<sup>st</sup> February, 2012.

How to cite this article: Banwat M.E., Yakubu N.W., Olalude E.O., Ogunsakin J.A. (2014). An assessment of the nutritional knowledge, practice and status of adult HIV/AIDS patients attending an art centre in Jos, north central Nigeria. *Afr. J. Food Sci. Technol.* 5(7):156-162