



International Research Journal of Plant Science (ISSN: 2141-5447)  
Vol. 14(2) pp. 01-5, April, 2023  
DOI: <http://dx.doi.org/10.14303/irjps.2023.08>  
Available online @ <https://www.interestjournals.org/plant-science.html>  
Copyright ©2023 International Research Journals

*Research Article*

# Effect of Grape Seed Extract (*Vitis Vinifera*) on Alternations of Ach and Ache Activities in Memory Defected Male Albino Rats

V.Uday Kiran<sup>1\*</sup>, M .Muniya Naik<sup>2</sup>, D.Veera Nagendra Kumar<sup>3</sup>, M.Rama Mohan<sup>4</sup>, A.Jayasankar<sup>5</sup>, P.Subramanyam<sup>6</sup>, P.Venkata Viswa Prasad<sup>7</sup>, Venkata Mohan Reddy<sup>8</sup>

<sup>1</sup>Department of Zoology, Loyola Degree College (YSRR) Pulivendula, AP, India

<sup>2</sup>Department of Zoology, Govt. Degree College, Rayachoty, AP, India

<sup>3</sup>Department of Zoology, Govt. Degree College for Men (A)-Kadapa, AP, India

<sup>4</sup>Department of Zoology, SV College of Arts & Computer Sciences, Proddatur, AP, India

<sup>5</sup>Department of Zoology, S.V.C.R.Govt. Degree College, Palamaner, Chittoor, AP, India

<sup>6</sup>Department of Botany, SKR & SKR Degree College for Women(A)Kadapa, AP, India

<sup>7</sup>Department of Zoology Govt.Degree College for Women, Rayachoty, AP, India

<sup>8</sup>Department of Zoology KSD College Porumamilla, AP, India

E- mail: [vempati.uday6@gmail.com](mailto:vempati.uday6@gmail.com)

## Abstract

A great source of different polyphenols with powerful antioxidant and disease-prevention qualities is grape seed extract (VGSE), which comes from the vine species *Vitis Vinifera*. There is a lack of research about the health advantages of VGSE on DNA damage, protein damage, labile iron activity, and enzyme inhibitory effects, despite the fact that VGSE has a wide range of biological activities that have the potential to improve human health. By exerting anti-oxidative, anti-inflammatory, anti-acetylcholinesterase, and anti-amyloidogenic effects, grape-derived extracts are natural sources of polyphenols that may promote healthy brain ageing. The acetylcholinesterase enzyme, which is discussed in this article, is involved in the cessation of cholinergic transmission, and it has been demonstrated that Alzheimer disease increases this enzyme's activity. We focus on the processes behind the neuromodulating properties of polyphenolic extracts and chemicals obtained from grapes, particularly resveratrol, grape seed extract, and extracts from grape leaves. However, additional study is needed to determine the most potent medicinal extracts and chemicals, as well as their brain bioavailability.

**Keywords:** Polyphenols, VGSE, Memory impairment, Neuromodulating properties.

## INTRODUCTION

The central and peripheral nervous systems both use the choline and acetic acid ester acetylcholine as a nerve impulse transmitter. Acetylcholine serves as the main neurotransmitter of the parasympathetic nervous system,

a part of the autonomic nervous system that lowers heart rate, contracts smooth muscles, widens blood vessels and stimulates bodily secretions. Acetylcholine can either promote or inhibit a process, therefore it can have both excitatory and inhibitory effects. Acetylcholine-containing vesicles are found at the terminals of cholinergic neurons.

**Received:** 02-Apr-2023, Manuscript No. IRJPS-23-75721; **Editor assigned:** 05-Apr-2022, PreQC No. IRJPS-23-75721(PQ); **Reviewed:** 19-Apr-2023, QCNo.IRJPS-23-75721; **Revised:** 22-Apr-2023, Manuscript No. IRJPS-23-75721 (R); **Published:** 27-Apr-2023

**Citation:** V.Uday Kiran (2023). Effect of Grape Seed Extract (*Vitis Vinifera*) on Alternations of Ach and Ache Activities in Memory Defected Male Albino Rats. IRJPS. 14: 08.

When a motor neuron's terminal receives a nerve impulse from the peripheral nervous system, acetylcholine is released into the neuromuscular junction. (Britannica.com) Alzheimer, (1906).

Acetylcholinesterase is a crucial enzyme in the cholinergic nervous system (AChE). Many different types of neurons degenerate as AD develops, quite apart from the massive loss of forebrain cholinergic neurons, which is followed by a gradual decline in acetylcholine (Mara-Salud et al., 2011). AChE, which hydrolyses acetylcholine, and Choline Acetyltransferase (ChAT), which synthesises it, are both affected. One of the main tenets of therapy designed to cure the cholinergic shortage is the importance of cholinergic function in cognition Bagchi et al., (2002). AChE inhibitors (AChE-I), which improve cholinergic transmission but have negligible and transient therapeutic benefits, are the mainstay of current AD treatment (Mara-Salud et al., 2011). Even while AChE activity has largely decreased in the AD brain, this is still the case Francis et al., (1999).

Cholinergic insufficiency is the most severe and pervasive metabolic disturbance in diseases that cause memory loss Murray & Pizzorno, (1998). It is now possible to observe acetylcholinesterase activity in the brain in real time García-Ayllón et al., (2011). Acetylcholine, choline acetyltransferase, and acetylcholine levels are all reduced as a result. Piperidyl derivatives have been employed in positron emission tomography (PET) to evaluate cortical acetylcholinesterase activity, such as N-[11C] methylpiperidyl-4-acetate ([11C] MP4A or [11C] PMP) 3 and [11C] methylpiperidyl-4-propionate ([11C] MP4P or [11C] PMP)3 ([11C]MP4A). Mild cognitive impairment is a result of early Memory Defect Disease and ageing naturally Hua et al., (2007). Only a tiny fraction of people with moderate cognitive impairment have memory issues by definition, but their general cognitive function and capacity to do daily tasks are unchange Goedert et al., (1989) d . But nothing is known about the role of the cholinergic system in mild cognitive impairment Lininger et al., (1998). We were interested in learning whether early memory impairment could be identified by changes to the cholinergic system in the temporal regions. (J O Rinne et al., 2003).

## MATERIALS AND METHODS

The present study was focused on the evaluation of the Cholinergic protective effect of Grape Seed extract on Memory defected rats Brain tissue Natella et al., (2002).

### Procurement and Maintenance of Experimental Animals

Healthy Wistar strain Albino rats, *Rattus norvegicus* of the same age group of 3 months, weighing  $160 \pm 20$  grams,

obtained from Sri Venkateswara enterprises, Bangalore were used as the experimental model in the present investigation Perry et al., (1991). Prior to experimentation, the rats were acclimatized according to the instructions given by Behringer (1973). They were housed in polypropylene cages under the controlled conditions of  $28 \pm 2^\circ\text{C}$  temperature with photoperiod of 12 hours light and 12 hours dark and 75% relative humidity maintained in the animal house of the Department Zoology, according to the ethical guidelines for animal protection and welfare bearing the Resolution No. 04/(i)/a/CPCSEA/ IAEC/ SVU/ KY- KPR / Dt. 28-03-2011. The rats were fed with standard pellet diet supplied by Sri Venkateswara Enterprises, Bangalore and water *ad libitum* throughout the period of experimentation.

### Preparation of Grape Seed Extract

Grape, as large clusters with red berries, was bought from a local fruit market in Tirupati, Pulivendula and Bangalore (Devanahalli) as *vitisvinifera*(Linn). Grape seeds were removed from the grapes, air dried (in shade) for one week and milled to fine powder (a particle size of  $< 0.4\text{mm}$ ). The grape seed powder was macerated in 75% ethanol for 72h at room temperature Rinne et al., (2003). The ethanol extract evaporated to remove ethanol, and grape seed extract was obtained as a lyophilized powder (Alireza Sarkaki et al., 2007). The resulting ethanolic crude extract was air dried and used in the present study Sarkaki et al., (2007).

### Administration of Tested Substance

Grape Seed Extract (GSE) 100 mg/kg body weight was dissolved in distilled water and given to the rat. A gavage tube was used to deliver the substance by oral route, which is clinically expected route for administration of GSE. The volume of administration was kept at 0.2 ml to the animal White & Foster, (2003).

**Grouping of Animals:** After the rats were acclimated to the laboratory conditions for 10 days before the experimentation, they were randomly divided into four groups. Each main group was again divided in to 2 sub-groups of six each and were housed in separate cages. These different groups of rats except control were treated with selected doses of Red grape seed ethanol extract and D-Gal as given below. Keeping in view the altered activity of rats during the nights compared to day time, all doses were given once in the morning hours in between 8 A.M. to 9 A.M (Table 1).

In the present study the experimental duration selected was 60days. D-Gal was given for first 30 days period to observe AD symptoms with the assessment of cognitive skills in rats (AD group). Further AD induced rats were again treated with D-Gal as well as Red grape seed ethanol extract simultaneously.

## RESULTS

From the results of the present research study, it was noticed that the Grape Seed Extract has significantly affected the total proteins and Cholinergic Neurotransmitters viz., ACh content and the AChE activity in brain tissue of D-Gal treated rats.

**Total Proteins:** Comparing experimental groups to control groups revealed significant differences in the amounts of total proteins; the AD-induced rats treated with D-Gal had the highest levels of protein. When compared to the AD-induced group, the AD-induced group treated with grape seed extract shown a notable increase in the level of total protein (Figure 1).


**Acetylcholine (ACh) Content:** The ACh content levels in all of the experimental groups significantly differed from

control rats and the AD-induced rats treated with grape seed extract had the highest ACh content. In rats treated with grape seed extract alone, a substantial rise was seen. The amount of ACh significantly decreased in AD-induced rats. When compared to the AD-induced group, the AD-induced group treated with Grape Seed Extract (GSE) concurrently shown a notable rise in ACh content in the experimental group (Figure 2).


**Acetylcholinesterase (AChE):** In contrast to ACh content, AD-induced rats showed higher AChE maximum activity levels than any other experimental groups. Rats administered with AChE Grape Seed Extract had considerably less activity overall when compared to the control group. The AD rats given with grape seed extract, however, showed the greatest inhibition Zhang et al., (2006). On the other hand,

**Table 1:** Grouping of animals.


<b>Group-I</b> (Control)	Control Rat
<b>Group-II</b> (AD)	Rat, Intraperitoneally(IP) administered with D-Gal (120 mg/kg body weight) up to end of the experiment (1 <sup>st</sup> day to 90 <sup>th</sup> day)
<b>Group-III</b> (GSE)	Rat, orally administered with Red grape seed ethanol extract (100mg/kg body weight) for 30 days.
<b>Group-IV</b> (AD+GSE)	Rat, Intraperitoneally injected with D-Gal (120 mg/kg body weight) once daily for first 30 days. From 31 <sup>st</sup> day onwards rats were administered with Red grape seed ethanol extract (100mg/kg body weight) for 30 days.



**Figure 1.** Graphical representation of Total Protein changes in rat brain tissue in control and experimental groups of rats.



**Figure 2.** Graphical representation of changes in Acetylcholine content ( $\mu$ moles of ACh/gm) in control and experimental groups of rats' brain tissue.



**Figure 3.** Graphical representation of changes in Acetylcholinesterase activity levels ( $\mu$ moles of Ach hydrolyzed/mg protein/h) in brain tissue of control and experimental groups of rats.

AD-induced rats had significantly higher AChE activity with the highest percent of elevation. The AChE activity was, however, recovered to nearly normal levels during the course of the trial in AD-induced rats treated with grape seed extract (Figure 3).

## DISCUSSION

The cholinergic system in rat brain tissue treated with Red Grape Seed Extract and D-Gal underwent considerable modifications in the current investigation. According to the findings, oral treatment of grape seed extract effectively preserved total protein levels and the activity of the cholinergic neurotransmitters ACh and AChE in experimental rats. It is thought that AChE's biological function is to stop impulse transmission by hydrolyzing the neurotransmitter ACh into acetic acid and choline (Nachmansohan and Neumann, 1975). By hydrolyzing the excitatory transmitter, Ach, AChE is a crucial regulatory enzyme that regulates the propagation of nerve impulses across cholinergic synapses (Milatovic et al., 2006).

Acetylcholine was the first neurotransmitter malfunction in Alzheimer's disease to be identified (ACh). It was thought that cholinergic dysfunction, which is necessary for short-term memory function, also contributed significantly to the short-term memory loss in AD (Francis et al., 1999). In the cortex and hippocampus, regions of the brain important in cognition and memory, markers for cholinergic neurons such as choline acetyltransferase and Acetylcholinesterase, enzymes responsible for synthesis and breakdown of ACh, respectively, are diminished (Francis et al., 1999). Cholinergic neurons are preferentially damaged in the nucleus basalis and the entorhinal cortex, where the early loss of neurons in AD patients occurs. Up to 90% of the cholinergic neurons in the nucleus basalis of Mynert may die as the disease worsens. (Wright et al., 1993).

Changes in the basal forebrain cholinergic system, notably in the hippocampus and cerebral cortex, have been linked to Alzheimer's Disease (AD) pathology. Normal ageing populations also experience cholinergic and memory deficiencies, however these dysfunctions are different from those seen with AD (Greig et al., 2005; Niewiadomska et al., 2009). When compared to the rats in the control group, the protein content in the AD-induced animals was considerably higher. This is a result of beta amyloid and tau protein production in the AD-affected brain. On treatment with GSE, levels of protein content were seen to be decreased, while in the AD+GSE experimental group of rats, levels of protein content were seen to be moderately elevated. Red Grape Seed Extract treatment, given concurrently, might return levels of protein content to normal. In Alzheimer's patients, choline acetyltransferase activity was found to be noticeably reduced. Given that  $\beta$ -amyloid has been demonstrated to inhibit choline absorption and ACh release in vitro (Zhong et al., 2009), cholinergic neurotransmission may be a particular target for  $\beta$ -amyloid.

The cholinergic system is crucial to memory and learning. Alzheimer's disease is linked with the loss of cholinergic neurons and decreased choline-acetyltransferase activity. Given that  $\beta$ -amyloid has been found to decrease choline absorption and ACh release in vitro, cholinergic neurotransmission may be a particular target for this substance. A number of negative behavioural symptoms associated with AD may potentially be influenced by changes in the central cholinergic system (i.e., depression, aggressive behavior, psychosis, and over activity). Cholinergic abnormalities have been observed in association with neurodegenerative conditions other than AD such as Parkinson's disease (Perry et al., 1999 & Furey et al., 2000).

## CONCLUSION

From the above results on brain cholinergic neurotransmitter system, it was finally concluded that, even though intake of grape Seed Extract for short term period improves all cholinergic neurotransmitters in ageing and delays the onset of Memory disfunctions.

The benefits and downsides of herbal plants can be explored in a variety of ways. Herbal drug has unwelcome constraints similar as a lack of unequivocal restrictions, a lack of impact, and a lack of substantiation. And, as can be seen, serological tests are employed in a wide range of operations, from detecting the presence of a specific disease to determining how to treat pathogens.

## ACKNOWLEDGEMENTS

I am grateful that my research work has received the required support from the research supervisor. I want to express my gratitude to my collaborators for all of their help in w this work. For giving computer lab space so that I could do this research, I also thank the Coordinator, DBT-supported Bioinformatics Infrastructure Facility, Department of Zoology, Sri Venkateswara University, Tirupati.

## REFERENCES

- Alzheimer, A. (1906). Über einen eigenartigen schweren Krankheitsprozess der Hirnrinde. *Zentralblatt für Nervenkrankheiten*. 25: 1134.
- Bagchi, D., Bagchi, M., Stohs, S.J., Ray, S.D., Sen, C.K. et al., (2002). Cellular protection with proanthocyanidins derived from grape seeds. *Ann N Y Acad. Sci.* 957: 260-270.
- Francis, P.T., Palmer, A.M., Snape, M., & Wilcock, G. K. (1999). The cholinergic hypothesis of Alzheimer's disease: a review of progress. *J Neurol*. 66: 137-147.
- García-Ayllón, M.S., Small, D.H., Avila, J., & Sáez-Valero, J. (2011). Revisiting the role of acetylcholinesterase in Alzheimer's disease: cross-talk with P-tau and  $\beta$ -amyloid. *Front Mol Neurosci*. 4: 22.
- Goedert, M.G.S.M., Spillantini, M.G., Jakes, R., Rutherford, D., & Crowther, R.A. (1989). Multiple isoforms of human microtubule-associated protein tau: sequences and localization in neurofibrillary tangles of Alzheimer's disease. *Neuron*. 3: 519-526.
- Hua, X., Lei, M., Zhang, Y., Ding, J., Han, Q., et al., (2007). Long-term D-galactose injection combined with ovariectomy serves as a new rodent model for Alzheimer's disease. *Life sci*. 80(20), 1897-1905.
- Lininger, S., Wright, J., Austin, S., Brown, D., & Gaby, A. (1998). *The Natural Pharmacy, Proanthocyanidins*. Virtuaol Health LLC. 199-200.
- Milatovic, D., & Dettbarn, W.D. (1996). Modification of acetylcholinesterase during adaptation to chronic, subacute paraoxon application in rat. *Toxicol Appl Pharmacol*. 136: 20-28.
- Murray, M.T., & Pizzorno, J.E. (1998). *Encyclopedia of natural medicine*. Three Rivers Press.
- Natella, F., Bellelli, F., Gentili, V., Ursini, F., & Scaccini, C. (2002). Grape seed proanthocyanidins prevent plasma postprandial oxidative stress in humans. *J Agric Food Chem*. 50: 7720-7725.
- Perry, G., Kawai, M., Tabaton, M., Onorato, M., Mulvihill, P., et al., (1991). Neuropil threads of Alzheimer's disease show a marked alteration of the normal cytoskeleton. *J Neurosci*. 11: 1748-1755.
- Rinne, J.O., Kaasinen, V., Järvenpää, T., Nägren, K., Roivainen, A., et al., (2003). Brain acetylcholinesterase activity in mild cognitive impairment and early Alzheimer's disease. *J Neurol*. 74: 113-115.
- Sarkaki, A., Farbood, Y., & Badavi, M. (2007). The effect of grape seed extract (GSE) on spatial memory in aged male rats. *Pak J Med Sci*. 23: 561.
- White, L.B., & Foster, S. (2003). The herbal drugstore: the best natural alternatives to over-the-counter and prescription medicines!. *Rodale*.
- Wright, C.I., Geula, C., & Mesulam, M.M. (1993). Neuroglial cholinesterases in the normal brain and in Alzheimer's disease: relationship to plaques, tangles, and patterns of selective vulnerability. *Annals of Neurology: Official J Am Neurol Ass Child Neurol Soc*. 34: 373-384.
- Zhang, Z.X., Zahner, G.E., Román, G.C., Liu, X.H., Wu, C.B., et al., (2006). Socio-demographic variation of dementia subtypes in china: Methodology and results of a prevalence study in Beijing, Chengdu, Shanghai, and Xian. *Neuroepidemiol*. 27: 177-187.
- Zhong, S.Z., Ge, Q.H., Qu, R., Li, Q., & Ma, S.P. (2009). Paeonol attenuates neurotoxicity and ameliorates cognitive impairment induced by d-galactose in ICR mice. *Neuro Sci*. 277: 58-64.