

Full Length Research Paper

Effect of low temperature on physico-chemical properties of different strawberry cultivars

Fardin Mirahmadi ^a, Qorban Mohammad Hanafi ^a, Mohammad Alizadeh ^b, Hamid Mohamadi ^a and Mohamad Sarsaifee ^c

^a Department of Food Science and Technology, Faculty of Agriculture, Azad University, Branch of Sanandaj, Sanandaj, Iran

^b Department of Food Science and Technology, Faculty of Agriculture, Urmia University, Urmia, Iran

^c Agricultural and Natural Resources Research Center of Kurdistan, Sanandaj, Iran

Accepted 24 January, 2011

Texture, anthocyanin content, pH, acidity, total soluble solids, total sugar, and sensory score of five strawberry cultivars (Kurdistan, Selva, Camarosa, Paroz, and Queen), stored at 6 °C, were investigated. The data indicated physical and chemical properties in all samples depended on cultivar during storage time since there were some contrasting responses among cultivars. 'Kurdistan' had the highest moisture content and showed decay symptoms previously. Except for 'Camarosa', TSS and total sugar increased in all cultivars. There were differences between cultivars concerning changes of acidity or pH over the storage period. In terms of flavor, 'Kurdistan', 'Camarosa', and 'Selva' cultivars were found better than the others at first day of cool storage. But 'Camarosa' and 'Selva' had a favorite colour among cultivars. Sign of fungal decay was observed in 'Kurdistan' samples at 7th day. This was the first study conducted on different cultivars of strawberry in our region.

Keywords: strawberry, physico-chemical properties, cool storage, sensory score.

INTRODUCTION

Strawberry is a non-climacteric fruit with limited shelf life due to its high sensitivity to chemical and microbial deterioration during postharvest storage and handling. It must be harvested at full maturity to achieve the maximum quality in relation to flavor and colour. As a consequence, strawberries should be harvested ready for consumption and due to this; strawberry fruit has a very short shelf-life at its best quality. In order to slow down metabolism and reduce deterioration prior to transport or storage, strawberries must be cooled to low temperatures after harvest. The postharvest life of strawberries can be extended by several techniques combined with refrigeration. Low temperatures and modified atmospheres are the tools for avoiding, at least partially, mold growth and fruit senescence, and thus extending the shelf-life of strawberries (Manning, 1996). One of the main factors in determining postharvest fruit deterioration

is expressive of softening, which reduces shelf life, facilitates pathogen infection and limits transportation and storage.

In spite of the wide use of low temperatures in strawberry preservation, not much is known about the effects of cool storage on chemical composition, in particular nutrients. When strawberry fruits are stored at low temperature, their shelf-life can be extended to at least one week. Nevertheless, the delay between harvest and storage at the proper temperature is critical for the success of the treatment. It was observed that fruits stored at low temperature, 6 h after harvest, showed undesirable changes in colour and texture and also a reduction of around 50% of water content in comparison to those that were immediately cooled after harvest (Nunes *et al.*, 1995). The main characteristics related to the quality of ripe strawberry fruit are texture, flavor (soluble sugars and organic acids) and colour (anthocyanin content). Change in texture is a consequence of the natural process of senescence and also of the atmosphere in which the fruit is stored.

*Corresponding author Email: m.xebat@gmail.com ; Tel.: +98 - 871- 32 84 932; fax: +98 -871- 32 84 931.

Besides the obvious changes in appearance, mold contamination can also promote undesirable changes in texture and contribute to reduced strawberry shelf-life.

Iran has produced 38500 tonnes of strawberry in 2007 (FAO, 2009). Kurdistan province grows about 80% of the total strawberry production in the Iran. In Kurdistan, the predominant cultivar used in planting and processing is 'Kurdistan'. The overall production in Kurdistan accounts for 30,000 tonnes. Processed strawberries account for about 4.5 tonnes in Kurdistan. Strawberry harvest begins in Kurdistan in May and peaks between May and June and continues until July. The objective of this work was to study texture, chemical composition and sensory score of five strawberry cultivars during one week of cool storage. The temperature of 6 °C was used because it is the average temperature usually applied to keep the fruit, when available for the final consumers.

MATERIAL AND METHODS

Origin and treatment of the Strawberries

The study was carried out on strawberry cultivars of Kurdistan, Camarosa, Selva, Queen and Paroz obtained directly from "Agricultural and Natural Resources Research Center of Kurdistan" on the same day. They were transported to lab in transparent polystyrene containers with a capacity of 500 g and stored at 6 °C. The fruits were carefully selected to be uniform in appearance and free from physical damage and deterioration.

Quality parameters

Firmness of strawberry was determined with a penetrometer for fruit firmness testing (The Wagner FT 02 penetrometer – Italy). The maximum force to penetrate the fruit sample was recorded as firmness. Each result was the mean of 10 determinations and was expressed in Newton (N) (Hernández-Muñoz *et al.*, 2006).

After firmness analysis, strawberries were cut into small pieces and homogenized in a grinder. Total soluble solids percent (%TSS) was measured in the juice of ground strawberries using an Atago RX-2500 digital refractometer (Atago Co. Ltd., Tokyo, Japan) at 20 °C. A drop of the juice was placed on the lens and the reading was taken in degree Brix. Calibration was made with deionized water and the lens was carefully rinsed between samples.

The pH was recorded with pH meter (pH-526; WTW Measurement Systems, Wissenschaftlich-Technische Werkstätten GmbH, Wellhelm, Germany). In the case of titratable acidity, six grams of ground strawberry was suspended in 20 mL of distilled water and then filtered. The filtrate titrated against 0.1 N NaOH up to pH 8.1 and expressed as citric acid. Total sugar as total reducing sugar was determined according to Iran's Standard (ISIRI, 2007).

Anthocyanin content of strawberries was determined using spectrophotometric method. 2 g of ground strawberry sample mixed with 20 ml of acidified methanol(1% HCl) using a homogenizer and then centrifuged at 2000 g for 15 min. Anthocyanin content was estimated as pelargonidin 3-glucoside at 510 nm, using a molar absorptivity of 36000 L.cm⁻¹.mol⁻¹.

Sensory score of strawberry samples were evaluated by ten trained staff members of the Kurdistan General Department of Standards and Industrial Research. The panelists were asked to evaluate the flavor. A twenty point scale was used where 20 = excellent and 1 = extremely poor. Accuracy and precision were statistically analyzed.

Data analysis

The results were analyzed using one-way analyses of variance (ANOVA) with the statistical software of SPSS (SPSS Inc., Chicago, IL, USA). Differences between means were studied with Duncan's test and differences at $p < 0.05$ were considered to be significant.

RESULTS AND DISCUSSION

Anthocyanin Determination

Anthocyanin contents during the storage period are shown in Figure 1. Figure 1 shows that based on initial amount of anthocyanin, cultivars can be classified in three groups: cultivars with high level of anthocyanin (Camarosa and Selva), medium level of anthocyanin (Kurdistan) and low level of anthocyanin (Queen and Paroz). The initial content of anthocyanin was 41.95 mg/100 g in Camarosa and low in Selva, while the content of this pigment was significantly ($\alpha=0.05$) lower (below 15 mg/100g) in Queen and Paroz. At the end of 7 days, anthocyanin content of Kurdistan and Paroz was not significantly different ($p>0.05$).

Figure 1 also shows that profile of anthocyanin change with storage time was similar among four cultivars (Camarosa, Selva, Queen and Paroz) but profile of Kurdistan was different from other cultivars. Fluctuations in anthocyanin content can originate from different the composition of anthocyanins and chemical changes followed by changes in their maximum absorption wavelength.

A total of eight pelargonidin- and three cyanidin-based anthocyanins were found in 39 strawberry cultivars (Baker *et al.*, 1994). Pelargonidin-3-glucoside was the most prevalent pigment, ranging from 100% to 82%. Strawberry pigments are very unstable due to the following chemical changes: hydrolysis of unstable aglycones, degradation of intermediaries, formation of copigment complexes with flavonoids, and degradation due to polyphenoloxidase. These reactions can lead to degrade or cause bathochromic shifts in anthocyanin colour (Wesche-Ebeling & Montgomery, 1990; Perera & Baldwin, 2001).

Texture Determination

Strawberries show a continuous decrease of cell wall content during ripening (Perkins-Veazie, 1995; Rosli *et al.*, 2004). Moreover, there are multiple chemical changes that involve pectins and the cellulose-xyloglucan framework resulting in solubilization of polyuronides and hemicelluloses (Knee *et al.*, 1977) and loss of neutral sugars (Nogata *et al.*, 1996; Rosli *et al.*, 2004). Huber (1984) stated that strawberry softening may be caused by the action of several enzymes, and may be accompanied by a loss of calcium.

Changes in firmness value between different cultivars




Figure 1. Effect of cool storage at 6°C on anthocyanin content (mg/100g) of five strawberry cultivars.




Figure 2. Effect of cool storage at 6°C on texture (N) of five strawberry cultivars.

during 7 days of storage at 6 °C are shown in Figure 2. The values of texture, measured on the first day of cold storage, ranged from 2.63 N for Kurdistan to more than

9.54 N for the Queen. Along with difference in initial values of the texture, the trend of textural change with time was also different among cultivars. The firmness of




Figure 3. Effect of cool storage at 6°C on pH of five strawberry cultivars.

all cultivars at the end of storage time was significantly higher than first day ($\alpha=0.05$). Texture changes were more evident in Camarosa, Selva, Queen, and Paroz. Kurdistan cultivar had high moisture and less firmness than the other cultivars. Soft texture would shorten the shelf-life of strawberry since it would become prone to mechanical injury and to mold contamination. The better characteristic texture of the Camarosa, Selva, Queen and Paroz makes them the most suitable for cool storage. These cultivars would also be an interesting choice when handling the fruit, since they are more resistant to physical injury and abrasion.

pH and titrable acidity

The change in pH of strawberries as a function of storage time is shown in Figure 3. pH of all cultivars ranged between 3.41 and 3.72, values that were above the average for ripe strawberry, pH of 3.3 (Green, 1971). Kurdistan cultivar had a low pH than the other samples during cool storage. The percentages of titrable acidity (TA), in all studied cultivars were between 0.52 and 0.86 during the storage period (Figure 4). These values are close to those reported by Green (1971) ranging from 0.6

to 2.3%. Marked changes in citric acid content were observed during fruit storage at cool temperature. Except for Paroz, all cultivars had a higher acidity than initial period at the end of storage time. After the 7 days of storage time, acidity level between Camarosa, and Queen and between Kurdistan and Selva was not significantly different ($\alpha 0.05$).

Organic, non-volatile acids are the second most important component of strawberry flavor, after soluble sugars. The main compound accounting for titratable acid (TA) is citric acid, which is predominant (over 90%) in strawberry. These acids regulate the cellular pH and may influence the anthocyanin stability and, as a consequence, the colour of the fruit. However, there is little published information about changes of pH and TA content in strawberry fruit stored at low temperatures. According to Nunes *et al.* (1995), cultivars 'Chandler', 'Oso Grande' and 'Sweet Charlie', stored at 1 °C, showed no differences in pH, but TA was slightly lower after 1 week. Remarkable changes of pH and citric acid content were observed by Gil *et al.* (1997) in stored strawberries of 'Selva' cultivar at 5 °C under different concentrations of CO₂ for 10 days. However, it was concluded that the effect was probably a consequence of the CO₂ since the increase of pH (and the decrease of TA) was paralleled




Figure 4. Effect of cool storage at 6°C on acidity (%) of five strawberry cultivars.

by the increase in concentration of CO₂ in the atmosphere.

The results of our study (Figure 3 and 4) clearly indicate that changes in pH and citric acid content during storage can also depend on cultivar. This information is important if there is a correspondence between the observed change in composition and fruit taste, since organic acids are one of the main components of strawberry flavor.

Total soluble solids and total sugar

The initial values for TSS ranged from 8.5 to 15 on the first day of storage in all cultivars. Changes in the soluble solids content of strawberries with storage time are shown in table 1. The results showed that changes in TSS during storage time vigorously depend on cultivar. The total soluble solid of mature strawberries has been reported to decrease under cold storage as a result of respiration (Garcia *et al.*, 1998). By contrast, the results obtained from Hernández-Muñoz *et al.* (2006) indicated that TSS increased with time. They reported changes by the considerable loss of water by strawberries during storage at 20 °C. The change in TSS content was mainly higher in Queen during storage time. After the 7 days of

storage, TSS level between Camarosa and Selva was not significantly different ($\alpha=0.05$). Since sucrose is the primary source of glucose and fructose, the increased amount of these monosaccharides could account for the decrease in sucrose level. However, the TSS increase indicates that sucrose synthesis had taken place during cool-storage.

At day 1, the amount of total sugar was below 7g/100g for Selva, Kurdistan, Paroz and Queen, while Camarosa had value above 9.5g/100g. Table 1 show that the TS were changed during the cold storage in all cultivars. Throughout the storage period the TS value was increased in all of them and except of the Camarosa fruits, the final sugar contents were higher than that of first day. Since strawberry does not have starch to support soluble sugar synthesis after harvest, this increase may be a consequence of cell-wall degradation.

Sensory scores

Sensory property (flavor) of strawberry cultivars on 1 and 7 days are shown in Figure 5. The sensory score of Paroz, Queen and Kurdistan strawberry samples changed during storage with a different trend for all the




Figure 5. Effect of cool storage at 6°C on sensory score (flavor) of five strawberry cultivars. Data presented are the means of 10 assays after 1 and 7 days of cool storage.

Table 1. Total soluble solids (TSS) (%) and total sugar (TS) (g/100 g) during storage of five strawberries cultivars

Days	Paroz		Camarosa		Kurdistan		Selva		Queen	
	TSS	TS	TSS	TS	TSS	TS	TSS	TS	TSS	TS
1	10.9± 0.1	6.1± 0.13	11.93± 0.2	9.9± 0.11	8.68± 0.12	5± 0.15	10.7± 0.3	5.2± 0.9	14.32± 0.2	5.5± 0.12
3	10.7± 0.3	6.93± 0.1	11.92± 0.1	7.6± 0.13	8.7± 0.2	5.7± 0.13	12.1± 0.15	6.9± 0.1	10.32± 0.14	6.65± 0.1
5	9.41± 0.21	7.1± 0.1	12.13± 0.3	7.7± 0.12	8.46± 0.11	5.7± 0.16	9.4± 0.2	5.9± 0.1	11.41± 0.1	7.46± 0.3
7	11± 0.1	10.3± 0.2	11.74± 0.1	8.3± 0.1	9.8± 0.14	6.4± 0.11	11.84± 0.1	7± 0.2	14.67± 0.18	6± 0.21

treatments. After 7 days of storage, reduction was more noticeable in Kurdistan cultivar. The flavor evaluation results showed that sensory score of Camarosa and Selva did not affect significantly by storage time. The sensory score of these two cultivars the highest among studied cultivars.

CONCLUSIONS

Lowering the storage temperature is an effective way to extend the strawberry shelf-life, maintaining the fruits edible for additional days. However, temperature can also induce small changes in some of the quality parameters studied. The data obtained reveal the importance of the cultivar in question since there were some contrasting responses among the varieties. Also, the initial values for some of the parameters were clearly different, indicating that the cultivar of strawberry is the most important factor

in determining post-harvest quality and extended shelf-life. This idea is reinforced by the data related to fruits cultivated at the same time under the same growing conditions.

ACKNOWLEDGEMENTS

We are grateful to Kurdistan General Department of Standards and Industrial Research for its laboratory support.

REFERENCES

- Baker J, Bridle P, Bellworthy SJ (1994). Strawberry juice colour: a study of the quantitative and qualitative pigment composition of juice from 39 genotypes. *J. of the Sc. of Food and Agric.* 64: 31–37.
 FAO (2009). FAOSTAT, available at : <http://faostat.fao.org/site/567/>

- Desktop Default. [aspx?Page ID=567#ancor](#)
- Garcia MA, Martino MN, Zaritzky NE (1998). Plasticized starch based coatings to improve strawberry (*Fragaria x ananassa*) quality and stability. *J. of Agric. and Food Chem.* 46: 3758–3767.
- Gil MI, Holcroft DM, Kader AA (1997). Changes in strawberry anthocyanins and other polyphenols in response to carbon dioxide treatments. *J. of Agric. and Food Chem.* 45: 1662–1667.
- Green A (1971). Soft fruits. In A. C. Hulme (Ed.), *the biochemistry of fruits and their products*, 2 (pp. 375–410). London and New York: Academic Press.
- Hernández-Muñoz P, Almenar E, José Ocio M, Gavara R (2006). Effect of calcium dips and chitosan coatings on postharvest life of strawberries (*Fragaria x ananassa*). *Post. Bio. and Tech.* 39: 247–253.
- Huber DJ (1984). Strawberry fruit softening: The potential role of polyuronide and hemicellulose. *J. of Food Sc.* 49: 1310–1315.
- ISIRI, Institute of Standards and Industrial Research of Iran, (2007). *Fruit juices – Test methods*, No. 2685, Karaj, Iran.
- Knee M, Sargent JA, Osborne DJ (1977). Cell wall metabolism in developing strawberry fruits. *J. of Exp. Bot.* 28: 377–396.
- Manning K (1996). Soft fruits. In G.B. Seymour, J.E. Taylor, & G.A. Tucker (Eds.), *Biochemistry of fruit ripening* (pp. 347–377). London: Chapman & Hall.
- Nunes MCN, Brecht JK, Morais AM, Sargent SA (1995). Physical and chemical-quality characteristics of strawberry after storage are reduced by a short delay to cooling. *Post. Bio. and Tech.* 6: 17–28.
- Perera CO, Baldwin EA (2001). *Biochemistry of fruits and its implications on processing* (2nd ed.). In D. Arthey & P.R. Ashurst (Eds.), *Fruit Processing, Nutrition, Products, and Quality Management* (pp. 19–36). Gaithersburg: Aspen Publishers.
- Perkins-Veazie P. (1995). Growth and ripening of strawberry fruit. *Hort. Rev.* 17, 267–297.
- Rosli HG, Civello PM, Martinez GA (2004). Changes in cell wall composition of three *Fragaria x ananassa* cultivars with different softening rate during ripening. *Plant Physiol. Biochem.* 42: 823–831.
- Wesche-Ebeling P, Montgomery MW (1990). Strawberry polyphenoloxidase: Its role in anthocyanin degradation. *J. of Food Sc.* 55: 731–734.