



International Research Journal of Plant Science (ISSN: 2141-5447)  
Vol. 14(2) pp. 01-8, April, 2023  
DOI: <http://dx.doi.org/10.14303/irjps.2023.10>  
Available online @ <https://www.interestjournals.org/plant-science.html>  
Copyright ©2023 International Research Journals

*Research Article*

# Estimation of Total Phenolic and Flavonoid Contents and Evaluation of Antioxidant Activity of Different Parts of *Cassia angustifolia* Vahl

Rashmi Rolaniya\*, R.A. Sharma, Jyoti Meena

Department of Botany, Laboratory of Plant Physiology and Biochemistry, Rajasthan, India

E- mail: [rolaniarashmi12@gmail.com](mailto:rolaniarashmi12@gmail.com)

## Abstract

The Caesalpiniaceae family includes the traditional medicinal plant *Cassia angustifolia* Vahl. The species is credited with a number of bioactive components, including phenolics, flavonoids, glycosides, coumarins, saponins, terpenoids, and alkaloids, among others, that contribute to a wide spectrum of therapeutic characteristics. The goal of the current study was to investigate the effect of different climatic conditions on total phenolic and flavonoid contents as well as on the antioxidant potential of different parts (root, stem, and leaves) of *C. angustifolia*. The plants were collected from three different states of India- Rajasthan, Gujarat, and Tamilnadu. Results of the current study showed that the selected plant parts possess significant amounts of total phenols and flavonoids. Our research revealed that methanolic extracts of leaves, roots, and stems from Tamilnadu had the maximum antioxidant activity, followed by Rajasthan and Gujarat. Rajasthan also had the highest levels of phenolic and flavonoid content, which was then followed by Tamilnadu and Gujarat. The results of a one-way ANOVA analysis showed that total phenolic and total flavonoid contents as well as antioxidant potential of the selected plant parts from the various sites were not significantly different. As a result, these plant parts can be utilised in place of harmful synthetic antioxidants and can be employed as natural antioxidants like red wine, green tea, and chocolates etc.

**Keywords:** Antioxidant potential, Total flavonoid content, Total phenolic content, Climatic condition.

## INTRODUCTION

As sources of remedies, medicinal plants are frequently employed as complementary therapeutic instruments for the prevention or treatment of numerous ailments. Recent studies have looked into how phenolic components including flavonoids, phenolic acids, tannins, etc. are primarily responsible for the antioxidant activity of plant parts (Nagavani and Rao, 2010; Cartea et al., 2010; Kaur & Mondal, 2014). Phenolics are ubiquitous secondary metabolites in plants. They include about 8000 bioactive chemicals. Numerous biochemical activities, including antioxidant, anti-mutagenic, and anti-cancerous, are exhibited by phenols. They can also alter gene expression significantly

(Nakamura et al., 2003). More than 4000 flavonoids of a plant origin have been discovered thus far. High flavonoid consumption lowers the risk of cancer and cardiovascular disease (Tapiero et al., 2002; Pamulaparthi et al., 2016; Ahmed et al., 2016). Flavonoids are a broad range of secondary metabolites in plants that are polyphenolic phytochemicals with variable phenolic structures. They include flavones, flavanone, flavanols, flavonols, and flavanonols (Hamamatsu et al., 2004; Santos et al., 2011; Chua et al., 2011; Laghari et al., 2010). Vegetables, fruits, flowers, grains, barks, roots, and stems all contain these phytochemicals. According to Cushnie & Lamb (2005), flavonoids show important biological activities such as antibacterial, antioxidant, anti-cancer, and anti-inflammatory activity (Laghari et al., 2011).

**Received:** 04-Apr-2023, Manuscript No. IRJPS-23-76574; **Editor assigned:** 07-Apr-2022, PreQC No. IRJPS-23-76574(PQ); **Reviewed:** 21-Apr-2023, QCNo.IRJPS-23-76574; **Revised:** 24-Apr-2023, Manuscript No. IRJPS-23-76574 (R); **Published:** 28-Apr-2023

**Citation:** Rashmi Rolaniya (2023). Estimation of Total Phenolic and Flavonoid Contents and Evaluation of Antioxidant Activity of Different Parts of *Cassia angustifolia* Vahl. IRJPS. 14: 10.

For all living beings, to survive on this planet, oxygen is a must. About 5% of oxygen is univalently reduced into oxygen-derived free radicals (ROS) like Superoxide Anions (O<sub>2</sub><sup>-</sup>), Hydroxyl (-OH), and Nitric Oxide (NO), which damage cellular components and cause tissue injury through covalent binding, during the process of oxygen utilisation in a normal physiological and catabolic process (Katsube et al., 2006; Krishna et al., 2019). Pathological illnesses like ischemia, asthma, arthritis, inflammation, neurodegeneration, Parkinson's disease, mongolism, the ageing process, and maybe dementia can be caused due to accumulation of free radicals. When antioxidant capacity is exceeded in biological system, the production of reactive oxygen species (ROS) during metabolism and other processes is increased which result into oxidative stress (Zima et al., 2001). Most frequent free radicals formed from nitrogen include nitric oxide (NO<sup>\*</sup>), peroxy nitrite anion (ONOO<sup>\*</sup>), hydroxy radical, hydrogen radical, and super oxide anion (O<sub>2</sub><sup>\*</sup>) (Pamulaparathi et al., 2016; Nagendrappa, 2005). Antioxidants are compounds which are present in low concentrations as compared to oxidizable substances but they can delay or stop those molecules from oxidising. By interacting with free radicals and functioning as oxygen scavengers, foods rich in antioxidants reduce the risk of chronic diseases such neurodegenerative disorders, cardiovascular diseases, the ageing process, cataracts, cancer, brain dysfunction, and other age-related degenerative diseases. Many synthetic antioxidants, including Butylated Hydroxyl Toluene (BHT), Butylated Hydroxyl Anisole (BHA), and Propyl Gallate (PG), are used to slow down the oxidation process in order to meet the rising demand for antioxidants, but their safety has long been questioned due to their unfavourable side effects and potential health risks (Pamulaparathi et al., 2016; Ahmed et al., 2016; Alhawari et al., 2021). Many research studies have turned to natural antioxidants in an effort to locate sources of antioxidants that would be secure, efficient, and affordable (Mundhe et al., 2011; Krishna et al., 2019). Free radical scavengers found in herbal medicines are renowned for their medicinal properties (Hakiman & Maziah, 2009). To stop the oxidation of the vulnerable substrate, plants create a highly astonishing range of antioxidant substances, such as carotenoids, flavonoids, ascorbic acid, etc. (Kaur & Mondal, 2014). Thus, there is a need to research and create natural antioxidants that are more effective and have lesser negative effects (Dela Torre et al., 2017).

The Caesalpiniaceae family includes the traditional medicine plant *C. angustifolia* Vahl. It is sometimes referred to as Cassia senna or Senna makkai. *C. angustifolia* is a native of Yemen, Saudi Arabia, and Egypt. It is a rapidly growing shrub between 5 and 8 metres tall that is widely farmed in Pakistan and India's hot, dry regions for its fruit and leaves

(Laghari et al., 2011). In both the British and American pharmacopoeias, this plant is included (Bameri et al., 2013). As an anti-helminthic, the leaves and pods of *C. angustifolia* are used as a powdered decoction treating intestinal worms. Additionally, it is frequently used as an anti-pyretic for cholera, splenic enlargements, typhoid, anaemia, toxicity, and genotoxicity brought on by *Escherichia coli* (Laghari et al., 2011). The studies published in the literature and the widespread use of *C. angustifolia* as a folk remedy for a variety of diseases call for more investigation to identify the substances causing its bioactivities (Ahmed et al., 2016). It has long been used as a purgative in folk medicine and is a well-known medication in traditional medicine (Wu et al., 2009). The majority of the world's pharmacopoeias also list this plant (Siddique et al., 2010). It is valued as a medicine because of its cathartic effects and is especially helpful in chronic constipation. Traditional purgatives include the leaves and pods of *C. angustifolia*; the most significant purgative components are Sennosides A, B, C, D, emodin, isorhamnetin, and essential oil (Siddique and Anis, 2007). *C. angustifolia* is a plant that greatly contributes to the production of commercial medications and has been researched in numerous regions of the world for a variety of therapeutic preparations in various methods. Its leaves are also used as a safe laxative (Arya, 2003; Laghari et al., 2011).

There are very few studies on *C. angustifolia*'s in vitro antioxidant tests, despite the fact that many reports have been published on the species' other therapeutic characteristics. Hence in the present study, antioxidant activity along with total phenolic and flavonoid contents was evaluated in different parts (stem, root and leaves) of *Cassia angustifolia* Vahl and effect of climatic conditions were also observed on those parameters.

## MATERIALS AND METHODS

### Sample Collection and Processing

Different plant parts of *Cassia angustifolia* Vahl (stem, root, and leaves) were collected from Rajasthan, Gujarat, and Tamilnadu in India. After washing the samples, the various plant sections were separated. Those underwent a 10-day period of room-temperature shade drying. The dry components were crushed into a powder. The powdered samples were retained and stored separately in an airtight container for use in additional phytochemical analysis.

### Sample Extraction

Using methanol as a solvent, the dried plant components were extracted using the cold percolation method. 10 g of the dry powder was placed in a conical flask with 100 ml of methanol and shaken at 120 rpm for 24 hours in an orbital shaker. The extracts were evaporated under vacuum after 24

hours and then filtered using Whatman filter paper no. 1 to remove debris.

### Determination of Total Phenolic Contents in the Plant Extracts

The Folin-Ciocalteu method was used to determine TPC (the Total Phenolic Content) (Singleton et al., 1965; McDonald et al., 2001). The reaction mixture was made by combining 0.5 ml of methanolic extract solution (1 mg/ml), 2.5 ml of water-dissolved 10% Folin-reagent, Ciocalteu's and 2.5 ml of 7.5% NaHCO<sub>3</sub>. The mixture was let to stand at 45° C for 15 minutes, and the phenols were then identified using a spectrophotometric technique. At 765 nm of wavelength, the absorbance was measured. The samples were made in triplicate, and the mean value of absorbance was calculated. Methanol was used to prepare the blank simultaneously instead of the extract solution. Gallic acid in methanol standard solution, 100–1000 mg/L, was used to create the standard curve. The amount of gallic acid equivalent (mg of GAE/g of dry weight), a frequent benchmark compound, was used to express the total phenolic content.

### Determination of Total Flavonoid Concentrations in the Plant Extracts

Using a slightly modified version of the aluminium chloride spectrophotometric technique (Chang et al., 2002), the concentration of TFC (Total Flavonoid Content) was measured. Plant extracts (0.5 ml) were dissolved in 1.5 ml of methanol, 0.1 ml of 10% aluminium chloride, 0.1 ml of 1 M potassium acetate, and 2.8 ml of distilled water before being incubated at room temperature for 30 minutes. At 415 nm, the reaction mixture's absorbance was gauged. The mean value of absorbance value was calculated for each experiment in triplicate, and values were represented as mean standard deviation. The standard quercetin solution in methanol was used to create the standard curve. The extracts' overall flavonoid concentration was measured in milligram of quercetin equivalents per gram of dry weight.

### Extraction of Flavonoids

Following the established method of Subramanian and Nagarajan, flavonoids were extracted from various plant parts (1969). Soxhlet extraction was performed on 100 grams of finely ground plant material using 500 ml of hot 80% methanol over the course of 24 hours. The filtrate was again extracted using petroleum ether, ethyl ether, and ethyl acetate in that order. To guarantee full extraction, each procedure was repeated three times. Because it contained a lot of fatty compounds, the petroleum ether fraction was discarded, and the ethyl ether fraction (which contained free flavonoids) was recovered. The bound flavonoids in ethyl acetate fractions were investigated. Each fraction underwent a 2-hour hydrolysis in 7 percent H<sub>2</sub>SO<sub>4</sub>. After filtering the

resulting combination, ethyl acetate was once more extracted from the filtrate. The ethyl acetate extract was extracted after being rinsed with distilled water until neutral. Both the free flavonoids and the bound flavonoids fractions of ethyl acetate were vacuum-dried, weighed, and kept in glass vials at 40° C until needed.

### Determination of Antioxidant Activity

**DPPH radical scavenging activity:** Using the method described by Bhat and Karim (2009), the 1, 1 - Diphenyl 2 - Picryl - Hydrazil (DPPH) was used to assess the extracts' ability to scavenge DPPH radicals. Peel extract was aliquoted and combined with 3.9 ml of a 0.1 mM DPPH methanolic solution. After vigorous vortexing, the mixture was left in the dark for 30 minutes. At 515 nm, the absorbance was measured in comparison to a methanol blank.

In order to determine the IC<sub>50</sub> values, a linear graph of concentration vs. percent inhibition was created. Each sample's antioxidant activity was calculated using the inhibition curve and given as IC<sub>50</sub>, which is the concentration needed to inhibit the generation of DPPH radicals by 50%. BHT (Butylated Hydroxy Toluene) (2.52 0.02) and ascorbic acid (2.54 0.00) were utilised as references

### Peroxidase Assay (POX)

200 mg plant sample was homogenised in 10 ml of phosphate buffer, chilled for 20 minutes, and then centrifuged at 10,000 rpm. The enzyme extract was taken from the clear supernatant. With the following modifications, the activity was assessed using the methodology of Chance & Maehly (1955). In addition, 0.2 ml of water, 0.3 ml of pyrogallol, and 2.4 ml of phosphate buffer were added Ahmed, (2016). Immediately after adding 0.1 ml of enzyme extract, the absorbance at 420 nm was measured to assess how much purpurogallin had produced Alhawarri, (2021). The enzyme activity, measured in mMoles per minute per gram dry weight, was calculated using the extinction coefficient of 2.8 mM<sup>-1</sup>cm<sup>-1</sup>.

### Statistical Analysis

The average of three analyses with standard deviation was used to express all experimental data, which were performed in triplicate. Additionally, linear regression analysis was used to get the IC<sub>50</sub> values.

## RESULTS AND DISCUSSION

Methanolic extracts of *C. angustifolia*'s leaves, stem, and root which were collected from several locations in India were analysed for their total phenolic and flavonoid content. Since methanolic extracts were discovered to have significant levels of both TPC and TFC, studies on the antioxidant activity of these extracts have been conducted.

These studies have included the DPPH assay and the POX enzyme assay. Although there have been a few reports on the in vitro DPPH assays of *C. angustifolia*, no research has been done on the species' methanolic extracts too far. Phenolics are significant secondary plant metabolites with known medicinal applications as antioxidants, anticarcinogens, and antimutagens. They have a history of lowering cardiovascular risks. The leaves from Rajasthan had the highest level of total phenolic content, which was found in the methanolic extract ( $21.98 \pm 3.65$  mg GAE/gdw), followed by the leaves from Tamilnadu ( $20.98 \pm 1.65$  mg GAE/gdw) (Table 1; Figure 1). The climate or plant environment is important which causes the variance in phenolic concentrations in the distinct places.


The class of polyphenolic chemical known as flavonoids includes substances that have powerful bioactivities against cancer, inflammation, ulcers, hepatotoxicity, allergies, and viruses. They are strong antioxidants because their phenolic hydroxyl groups enable them to efficiently scavenge ROS. The leaves collected Rajasthan had the highest levels of flavonoids ( $38.86 \pm 11.09$  mg QE/gdw), followed by the leaves gathered in Tamilnadu ( $33.12 \pm 09.02$  mg QE/gdw) (Figure 2 & 3). The extraction method and the plant's environment both affect how much flavonoid is present overall in plant

extracts. High phenolic and flavonoid content in the extracts may have directly influenced the species' antioxidant activity. Given that plants with high flavonoid content can act as good sources of antioxidants to help increase an organism's overall antioxidant capacity and provide protection from lipid peroxidation, the presence of good amounts of phenolic and flavonoid contents demonstrated that both species are good sources of natural antioxidants (Sharififar et al., 2009).

The most popular technique for assessing the antioxidant activity of natural antioxidants is DPPH. According to Janaszewska et al. (2002), different antioxidants react with DPPH at varying rates, which affects how well they scavenge free radicals. Methanolic plant extracts may have scavenging potential since there are free hydroxyl groups available. Our findings also imply that there are more free hydroxyl groups in leaf methanolic extracts since stem extract has the highest DPPH free radical scavenging activity, followed by root and leaf extracts. The most effective plant parts for antioxidant potential were collected from Rajasthan (IC<sub>50</sub>=98.309.03 mg/L [Leaves], 285.910.04 mg/L [Stem], 109.506.02 mg/L [Root]), followed by Tamilnadu (IC<sub>50</sub>=100.108.01 mg/L [Leaves], 302.310.87 mg/L [Stem], 129.609.76 mg/L [Root]), and Gujarat (IC<sub>50</sub>= 111.4  $\pm$  10.05 mg/ml [Leaves],

**Table 1:** Total phenolic, total flavonoids, and isolated flavonoid contents in *Cassia angustifolia vahl*.

Plant part	Locations	Total phenolic content (mg GAE/gdw)	Total flavonoid content (mg QE/gdw)	Isolated flavonoid (mg/d.dw)		
				Free (F)	Bound (B)	Total (F+B)
Leaves	R	21.98 $\pm$ 3.65	38.86 $\pm$ 11.09	2.134	1.296	3.397
	G	19.98 $\pm$ 2.05	28.06 $\pm$ 10.01	1.156	0.978	2.285
	T	20.98 $\pm$ 1.65	33.12 $\pm$ 09.02	2.101	1.002	2.976
Stem	R	12.47 $\pm$ 8.84	23.87 $\pm$ 9.32	1.074	1.523	1.597
	G	10.47 $\pm$ 4.05	13.77 $\pm$ 8.10	0.037	0.987	0.972
	T	11.47 $\pm$ 6.36	19.34 $\pm$ 7.02	1.001	1.002	1.203
Root	R	19.25 $\pm$ 0.01	5.59 $\pm$ 2.58	2.384	1.341	3.345
	G	16.25 $\pm$ 0.03	3.12 $\pm$ 1.06	1.196	0.149	2.540
	T	17.25 $\pm$ 0.02	4.38 $\pm$ 2.08	2.003	1.011	3.009



**Figure 1.** Total Phenolic C

content (mgGAE/gdw) of *Cassia angustifolia* Vahl.




Figure 2. Total Flavonoid Content (mgQE/gdw) of *Cassia angustifolia* Vahl.




Figure 3. Isolated Flavonoids (mg/d.dwt) of *Cassia angustifolia* Vahl.

Table 2: Evaluation of antioxidant activity of plant parts by DPPH assay.


Name of plant part	Locations	IC <sub>50</sub> (mg/L)
		<i>Cassia angustifolia vahl.</i>
Leaves	R	98.3±09.03
	G	111.4±10.05
	T	100.1±08.01
Stem	R	285.9±10.04
	G	320.4±11.35
	T	302.3±10.87
Root	R	109.5±06.02
	G	135.7±09.25
	T	129.6±09.76

$320.4 \pm 11.35$  mg/L [Stem],  $135.7 \pm 09.25$  mg/L [Root]) (Table 2; Figure 4). Leaf part of all location found to be more potent antioxidant than root and stem. In general, the antioxidant activity will be higher the lower the IC<sub>50</sub> values. The discrepancy in their IC<sub>50</sub> values indicates that the scavenging activity of the methanolic extract of the root is less than that of the leaves. The different polarity and solvent extract dissolving capacities amongst them may be the cause of the variation in their scavenging abilities. By using a one-way ANOVA analysis, no discernible difference was detected between the IC<sub>50</sub> values of in similar plant parts collected from various locations however, in a single plant, different parts showed varied antioxidant potential significantly. The potential antioxidant activity of a substance is indicated by its capacity to reduce power. The quantity of reductions a substance has determines how much it can reduce. These reductions block the chain of free radicals by giving a hydrogen ion. Higher amounts of reductions were found in the extracts as evidenced by the reducing power ability of methanolic extracts, which rose as the concentration of the test extract increased. The irreparable harm caused by oxidative stress depends on the production of OH\* radicals. Hydroxyl radicals are produced by the Fenton reaction and

are very reactive in biological systems. You can calculate a test compound's ability to scavenge hydroxyl radicals by looking at how free radicals affect deoxyribose's ability to degrade (Subhashini et al., 2011).

The results of the DPPH assay were further corroborated by the peroxidase enzyme assay, which showed that leaves have more antioxidant potential than roots. More effective extracts came from Tamilnadu ( $0.246 \pm 0.022$  mMole/min/g.dw [Leaves],  $0.153 \pm 0.020$  mMole/min/g.dw [Stem],  $0.211 \pm 0.120$  mMole/min/g.dw [Root]), Rajasthan ( $0.198 \pm 0.019$  mMole/min/g.dw [Leaves],  $0.131 \pm 0.017$  mMole/min/g [Stem],  $0.203 \pm 0.119$  mMole/min/g.dw [Root]) and Gujarat ( $0.083 \pm 0.011$  mMole/min/g.dw [Leaves]  $0.076 \pm 0.010$  mMole/min/g.dw [Stem],  $0.108 \pm 0.108$  mMole/min/g.dw [Root]). Roots were discovered to be more powerful than leaves and stems among all plant parts (Table 3; Figure 5).

As a result, we can conclude that in the current study, methanolic extracts of leaves, roots, and stems collected from Tamilnadu showed more potent antioxidant activity than those from Rajasthan and Gujarat, while Rajasthan had the highest levels of phenolic and flavonoid content, followed by Tamilnadu and Gujarat



**Figure 4.** Evaluation of antioxidant activity of different parts of *Cassia angustifolia* vahl. by DPPH assay.

**Table 3:** Evaluation of antioxidant activity by peroxidase (POX) enzyme assay.

Name of plant part	Locations	Peroxidase activity (mMole/min/g.dw)
		<i>Cassia angustifolia</i> vahl.
		Methanolic extract
Leaves	R	$0.198 \pm 0.019$
	G	$0.083 \pm 0.011$
	T	$0.246 \pm 0.022$
Stem	R	$0.131 \pm 0.017$
	G	$0.076 \pm 0.010$
	T	$0.153 \pm 0.020$
Root	R	$0.203 \pm 0.119$
	G	$0.108 \pm 0.108$
	T	$0.211 \pm 0.120$




Figure 5. Evaluation of peroxidase enzyme activity of different parts of *Cassia angustifolia* vahl.

## CONCLUSION

The current study demonstrates that different parts of *Cassia angustifolia* Vahl have significant flavonoid and phenolic content. The study's findings also indicated that these plant parts possess a significant amount of antioxidant compounds. Gujarat and Tamilnadu are followed in antioxidant activity and bioactive constituent richness in plant parts collected from Rajasthan. The results of the current study may help scientists and researchers in formulating plans for creating food crops high in antioxidants. Further research is needed to fractionate the extract, identify the bioactive compound(s), and conduct in vivo tests on its medicinal active ingredients in order to pinpoint their precise mechanisms of action, enhance nutritional benefits, and create high-quality natural pharmaceutical products.

## ACKNOWLEDGEMENT

Authors are grateful to the Head, Department of Botany, University of Rajasthan, Jaipur, Rajasthan, for providing the necessary resources for the research.

## REFERENCES

- Ahmed, S.I., Hayat, M.Q., Tahir, M., Mansoor, Q., Ismail, M., et al., (2016). Pharmacologically active flavonoids from the anticancer, antioxidant and antimicrobial extracts of *Cassia angustifolia* Vahl. *BMC Complementary Alternative Med.* 16: 1-9.
- Alhawarri, M.B., Dianita, R., Razak, K.N.A., Mohamad, S., Nogawa, T., et al., (2021). Antioxidant, anti-inflammatory, and inhibition of acetylcholinesterase potentials of *Cassia timoriensis* DC. flowers. *Molecules.* 26: 2594.
- Althman, M., Bhat, R., & Karim, A.A. (2009). Antioxidant capacity and phenolic content of selected tropical fruits from Malaysia, extracted with different solvents. *Food Chem.* 115: 785-788.
- Arya, R. (2003). Yield of *Cassia angustifolia* in combination with different tree species in a silvi-herbal trial under hot arid conditions in India. *Bioresour Technol.* 86: 165-169.
- Bameri, Z., Amini-Boroujeni, N., Saeidi, S., & Bazi, S. (2013). Antibacterial activity of *Cassia angustifolia* extract against some human pathogenic bacteria. *J Nov Appl Sci.* 2: 584-586.
- Cartea, M.E., Francisco, M., Lema, M., Soengas, P., & Velasco, P. (2010). Resistance of cabbage (*Brassica oleracea capitata* group) crops to *Mamestra brassicae*. *J Econ Entomol.* 103: 1866-1874.
- Chance, B., & Maehly, A. C. (1955). [136] Assay of catalases and peroxidases.
- Chang, C.C., Yang, M.H., Wen, H.M., & Chern, J.C. (2002). Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *J Food Drug Anal.* 10.
- Chua, L.S., Latiff, N.A., Lee, S.Y., Lee, C.T., Sarmidi, M.R., et al., (2011). Flavonoids and phenolic acids from *Labisia pumila* (Kacip Fatimah). *Food Chemis.* 127: 1186-1192.
- Cushnie, T.T., & Lamb, A.J. (2005). Antimicrobial activity of flavonoids. *Int J Antimicrob Agents.* 26: 343-356.
- Hakiman, M., & Maziah, M. (2009). Non enzymatic and enzymatic antioxidant activities in aqueous extract of different *Ficus deltoidea* accessions. *J Med Plant Res.* 3: 120-131.
- Hamamatsu, S., Yabe, K., & Nawa, Y. (2007). Compositions of anthocyanin and other flavonoids in cultured cells of rabbiteye blueberry (*Vaccinium ashei* Reade cv. Tifblue). *Food Sci Technol Res.* 10: 239-246.
- Katsube, T., Imawaka, N., Kawano, Y., Yamazaki, Y., Shiwaku, K., et al., (2006). Antioxidant flavonol glycosides in mulberry (*Morus alba* L.) leaves isolated based on LDL antioxidant activity. *Food Chem.* 97: 25-31.
- Kaur, S., & Mondal, P. (2014). Study of total phenolic and flavonoid content, antioxidant activity and antimicrobial properties of medicinal plants. *J Microbiol Exp.* 1: 00005.
- Krishna, K.V.V.S., Venkatesh, B., Madhavi, A., Devi, B. V., Jaldi, C.E., et al., (2019). Evaluation of Antioxidant Activity of Panchachurnam: An Ayurvedic Polyherbomineral Formulation. *Res J Pharm Technol.* 12: 4841-4847.
- Laghari, A.H., Memon, S., Nelofar, A., Khan, K.M., Yasmin, A., et al., (2010). A new flavanone with urease-inhibition activity isolated from roots of manna plant camelthorn (*Alhagi maurorum*). *J Mol Struct.* 965: 65-67.

- Laghari, A.Q., Memon, S., Nelofar, A., & Laghari, A. H. (2011). Extraction, identification and antioxidative properties of the flavonoid-rich fractions from leaves and flowers of *Cassia angustifolia*. *Am J Anal Chem.* 2: 871.
- Torre, G.D., Arollado, E.C., Atienza, A.A., & Manalo, R.A.M. (2017). Evaluation of antioxidant capacity and identification of bioactive compounds of crude methanol extracts of *Caesalpinia pulcherrima* (L.) Swartz. *Indian J Pharm Sci.* 79: 113-123.
- Wu, Q.P., Wang, Z.J., Tang, L.Y., Fu, M.H., & He, Y. (2009). A new flavonoid glucoside from *Cassia angustifolia*. *Chin Chem Lett.* 20: 320-321.
- Zima, T., Fialová, L., Mestek, O., Janebová, M., Crkovská, J., et al., (2001). Oxidative stress, metabolism of ethanol and alcohol-related diseases. *J Biomed Sci.* 8: 59-70.