

Review

Fatigue experience among cancer patients receiving chemotherapy

¹Bilal S.H Badr Naga and ^{2*}Manal I. Kassab

¹Oncology Nurse Specialist, Master Student, Hashemite University, Jordan
^{2*}Faculty of Nursing, Jordan University of Science and Technology (JUST), Jordan
Associate (Clinical Fellow) in Nursing at University of Technology, Sydney (UTS)

Abstract

A systematic review conducted between January 2003 and October 2010 to describe what is known about strategies used to prevent Cancer-related fatigue (CRF); to determine factors that have been associated with treatment of cancer and to identify the risk factors associated with methods of treatment of cancer that causes fatigue. Pub Med database and reference lists of identified articles were used. Variations in research design and methods including sample characteristics, interventions and measurements used to reduce fatigue were recorded. Ten RCTs out of 20 studies were included. Eight studies showed that cancer patients experience fatigue during and post chemotherapy besides having physical, social, and psychological problems. Two studies found that patient using steroids has a significant increase in fatigue; while, two studies reported that CRF associated with lower quality of life regardless of chemotherapy regimens. Eight of ten studies showed regular committed exercise (walking or swimming) resulted in less fatigue among patient participating in exercise programs. Evidence from reviewed studies supports the inclusion of physical and psychological support for managing CRF and they are considered important factors for predicting changes in patients' quality of life. They should be included in care plan of patients undergoing chemotherapy. Successful symptom management for patients can help maintain effective chemotherapy, physical/social wellbeing, and reduce emotional distress of patients. There is need to expanded nursing educational programmes regarding CRF assessment and possible management options to reduce severity of CRF and improve their quality of life.

Keywords: Cancer-Related Fatigue, Quality of Life, Chemotherapy, Jordan.

INTRODUCTION

Fatigue is one of the most common side-effects of cancer and its treatments (Hofmana et al., 2007, Prue, Rankin, Allen et al., 2006). Results of a multi-centre patient survey revealed that cancer patients identify fatigue as an important problem which affects their daily activities for more of the time than either nausea/vomiting or cancer pain. CRF is usually characterized as an overall lack of energy, cognitive impairment, somnolence, mood disturbance, or muscle weakness (The National

Comprehensive Cancer Network (NCCN), 2012). These symptoms occur with cancer and cancer therapy and are not relieved by rest or additional sleep and often interfere with daily activities.

Cancer-related fatigue that associated with chemotherapy has been associated with other symptoms, including pain, difficulty sleeping and muscle weakness (Cleeland, 2007). Cancer-related fatigue may interact with other common adverse effects of chemotherapy drugs, such as nausea and vomiting, by increasing their perceived severity. This could increase their impact on patient activity, and challenge patients' ability to complete their recommended treatment on the optimal schedule. In addition to having a significant influence on the quality of

*Corresponding Author Email: manal_kassab@yahoo.com

life (QOL) during chemotherapy, CRF may be present even before treatment begins; it can increase during the course chemotherapy; and it can persist at a higher-than-baseline rate, sometimes for years, after cancer treatment is finished (Nail and King (1987).

Cancer-related fatigue has been under-reported, under-diagnosed and under-treated (NCCN, 2012). Health care professionals have been challenged in their efforts to help patients manage this distressful symptom and to maintain the quality of patients' life. Compared with other health care providers, nurses spend most of their time with patients and their families (Dickinson, Clark, and Sque, 2008). Nurses play a major role in the care of individuals and their families in all stages of cancer, from diagnosis to death. Nurses deal frequently with cancer patients and trying to maintain high quality of care, alleviate suffering, decrease side effect and complications of cancer treatment (Dickinson, Clark, and Sque, 2008). Nurse's knowledge and experience about CRF can shape their attitudes toward care for cancer patients (William, Dale, Godley, and Neimeye, 2003). In addition, nurses are in the most immediate position to provide care, comfort and counseling for patients and families at the stage of cancer management (Dickinson, Clark, and Sque, 2008).

Background

Cancer-related fatigue (CRF) is a universal distressing symptom among cancer patients who are receiving cancer treatments (Gibson, et al., 2005). The National Comprehensive Cancer Network (NCCN) defined CRF as: "*distressing persistent, subjective sense of physical, emotional and/or cognitive tiredness or exhaustion related to cancer or cancer treatment that is not proportional to recent activity and interferes with usual functioning*" (NCCN, 2012, p.FT1). Cancer-related fatigue has been reported as the most important symptom that impairs the quality of life and daily patients' activities (Rayan, et al., 2007). According to NCCN, 70% to 100% of cancer patients experience CRF and most cancer patients suffer from CRF while receiving chemotherapy, radiation, or bone marrow transplant (NCCN, 2012). Cancer patients experience CRF resulting from the disease process, treatment modalities, psychological burdens, and worsen during chemotherapy course and persists for months after completing the treatment (Mustian, Morrow and Carroll, 2007).

A study conducted by Erickson, et al., (2010) aimed to explore factors affecting fatigue. The researchers recruited 20 adolescents with cancer receiving chemotherapy. Twelve cancer patients in the study reported brief peaks of moderate to severe CRF that

occurred one to four days after each week one of chemotherapy administration. In addition, the researchers found that the factors that made CRF worse included "not being able to eat before procedures, going to the bathroom a lot, getting chemotherapy in the morning, and just getting chemotherapy".

Alison et al., (2005) mentioned that patients with cancer have identified CRF as symptom that causes them major distress as they live with the disease and its treatment sequel. Many did experience physical, social, and psychological distress as a result of having fatigue.

Relationship between Chemotherapy and Fatigue

Wielgus et al., (2009) found that higher CRF is associated with chemotherapy treatment. Hofman and colleagues (2007) found that more than 80% of patients receiving chemotherapy report CRF as a significant side effect of treatment. At the same study they found that 88% of patients who had chemotherapy reported that CRF had affected their activities of daily living. Lucia and colleagues (2003) explain the relation between cancer and CRF and found that 70% of people with cancer report feelings of CRF during and after adjuvant chemotherapy.

Factors Associated with Cancer-Related Fatigue

Hwang, et al., (2003) in their study proposed a conceptual model with three dimensions (situational, biological, and psychological dimensions) that predict cancer-related fatigue. The situational dimension represents demographic information including age, gender, stage of cancer, active cancer treatment, and caregiver status. The biological dimension can be described by serum chemistry profile to identify anemia caused by chemotherapy or radiotherapy (Berndt, et al., 2005). The impact of anemia on CRF may be different depending on onset time, patient age, and comorbidity (Berndt, et al., 2005). Psychological factors, such as depression and anxiety, may contribute to the development of chronic CRF before and after chemotherapy among patients with solid tumors (Wasteson et al., 2009). Distress after a diagnosis of cancer can be caused by the initial fatigue, and other side effects of upset, like insomnia which may also increase in patients undergoing chemotherapy (Wasteson et al., 2009).

Furthermore, Yeh et al., (2008) investigated the relationships between clinical factors (including hemoglobin value, chemotherapeutic agents, and corticosteroid use) and changing patterns of CRF before

and for the next 10 days following the start of a new round of chemotherapy in children with cancer. The researchers found that CRF levels were changed significantly over time; patients have more problems with CRF in the first few days after the start of chemotherapy. Also, the researchers reported that Corticosteroid use and hemoglobin value were associated with significant increases in CRF that were sustained for several days and reached the highest level of CRF at day 5 for those receiving concurrent steroids. In addition,

Yeh et al., (2008) reported that the association of chemotherapeutic agents with CRF varied between patient self-report and parent report, but the type of chemotherapeutic agents used was not associated with most changes in fatigue. This study supports that lower hemoglobin level as a significant contributor to fatigue, thus, frequent hemoglobin check might increase the nurses' understanding of CRF that accompanies chemotherapy treatment.

Berger et al. (2009) examined the relationships among cancer-related fatigue, physical and mental quality of life (QOL) and different chemotherapy regimens in patients prior to, during, and after treatment. The researchers reported that CRF and mental (QOL) changed significantly over time for all regimens, but the patterns of change did not differ based on regimen. Physical (QOL) changed significantly over time for all regimens. Higher CRF was correlated with lower physical and mental (QOL) prior to and 30 days after the final treatment, regardless of regimen. The results of this study assure that higher CRF associated with lower (QOL) regardless of the chemotherapy regimens. Thus, study recommended that nurses should screen patients for CRF using a visual analog scale (VAS), assess for contributing factors and to integrate evidence-based CRF interventions as early as possible to reduce CRF and prevent lower quality of life during treatment.

Measures Used to Decrease Cancer-Related Fatigue

Yurtsever (2007) studied the experience of CRF in patients receiving chemotherapy. The researcher focused on measures taken by patients to cope with fatigue. Cancer-related fatigue was found to be affected by patients' in their daily activities, ages, gender, treatment, and symptoms related to the chemotherapy. Yurtsever (2007) found that the majority of cancer patients receiving chemotherapy (86%) experienced fatigue; and 73% stated that they coped with CRF by decreasing their activities and resting more. In addition, taking care with their nutrition (12%), exercising (5%), reading a book or newspaper (3%), listening to music (3%), drinking lots of fluids (3%), watching television (3%), trying to cope with

the pain (3%), and massage (3%). Additionally, when the patients used these measures, 26% stated that the measures were "partially effective" in decreasing their CRF and 37% stated that they were "ineffective." Age was not found to be a significant contributing factor that is affecting the level of fatigue. However, other related factor including gender, length of illness, number of chemotherapy courses, and patients' symptoms were found to have an effect on level of fatigue. The findings of Yurtsever may refer to the majority of patients experienced CRF coped with less activity, and more resting.

On the contrary, Kuchinski et al., (2009) did a systematic review to determine the effect of exercise in decreasing CRF for patients receiving chemotherapy and radiation. The researchers found that eight out of ten studies showed regular committed exercise (walking, bicycling or swimming) resulted in less CRF among patients participating in exercise programs. Participants who walked at least 60 minutes per week in three or more sessions demonstrated an increased functional capacity and activity level, with minimal increase in CRF compared to the control group.

In another study done by Blaney et al., (2010) who explored the barriers of using exercise among patients recently diagnosed with cancer and suffering of fatigue. The researchers reported that Cancer-related fatigue was associated with barriers such as physical problem, social isolation, and difficulty of making a routine exercise.

Lee et al., (2008) explored the relationship between fatigue, hemoglobin, and the coping strategies used by cancer patients receiving chemotherapy. The researchers found that majority of patients had a baseline hemoglobin level of 12 g/dl and a significantly greater mean cancer-related fatigue score than patients with hemoglobin >12 g/dl. Cancer-related fatigue levels were significantly higher in patients receiving a third course of chemotherapy than in those receiving first course. The most commonly used management strategy was energy conservation (sitting and lying down). However, participants rated exercise, sleep, going to bed early and walking as the most effective. Distraction techniques as listening to music, reading books and visiting with friends had low-to-moderate effectiveness. The most effective coping strategy was chatting with others; back massage and relaxation training which they were found to be moderately effective in reducing chemotherapy-induced fatigue. Thus, implementation of these coping strategies may prevent CRF and promote the quality of life.

Yeşilbalkan (2009) did a quasi-experimental, descriptive study to determine whether a nursing educational interventions decreased the perception of CRF among gastrointestinal (GI) cancer patients receiving chemotherapy for the first time. The researcher

assessed cancer-related quality of life by using Fatigue Inventory, Piper Fatigue Scale, and the European Organization for Research and Treatment of Cancer Quality of Life (EORTC QLQ C-30) scale before their first cycle of chemotherapy, on the 10th day after (T1), and again 10 days after the second cycle of chemotherapy (T2). Patients received an individual educational intervention at baseline, T1, and T2 based on the results of their CRF assessment. Patients were given an educational booklet on CRF prior to treatment and symptom specific booklets as required at T1 and T2. The researcher found that at baseline, patients generally reported moderate levels of fatigue for each subscale of the (PFS) (behavioral, affective, sensory, and cognitive), but the levels decreased with each subsequent intervention (i.e., at T1 and T2). Following the educational intervention, mean scores in the functional domain of the EORTC QLQ C-30 (physical, role, cognitive, emotional, social, and global QOL) have been increased while symptoms (fatigue, pain, lack of appetite) have been decreased at both T1 and T2, compared to baseline. These findings emphasized importance of developing nursing educational program regarding CRF assessment. This is crucial for the provision of appropriate educational interventions to patients prior to chemotherapy in order to help in reducing severity of CRF and improve quality of patient's life.

CONCLUSION

Cancer-related fatigue is the most commonly reported side effect of cancer and its associated treatment options, mainly chemotherapy. Cancer-related fatigue differs from that induced by other causes, such as sleep disturbance and exertion, as the latter are typically alleviated by a period of rest. This literature review considered the effect of several factors including exercise on CRF among patients treated with chemotherapy. Evidence from these studies supports the inclusion of scheduled exercise in the care plan of patients undergoing chemotherapy. Patients with cancer may be challenged to do exercise with nursing support. This support can be made more effective if the recommended exercise program is regularly adjusted to the patients' health status with consideration to other factors such as level of hemoglobin, age, course of chemotherapy. Nurses' awareness of the role of exercise in managing CRF related can provide better education that benefits patients. Results of included studies indicated that exercise and psychological interventions provided reductions in cancer-related fatigue. Specifically, multimodal exercise and walking programs, restorative approaches, supportive – expressive, and cognitive –

behavioural psychosocial interventions show promising potential for ameliorating cancer-related fatigue. With improved methodological approaches, further research in this area may soon provide clinicians with effective strategies for reducing CRF and enhancing the lives of cancer patients.

REFERENCES

- Andrykowski MA, Curran SL, Lightner R (1998). Off-Treatment Cancer-related fatigue in Breast Cancer Survivors: A controlled Comparison. *J. Behav. Med.*; 21:1–18
- Berger AM, Lockhart MK, Agrawal S (2009). Variability of Patterns of Cancer related fatigue and Quality of Life Over Time Based on Different Breast Cancer Adjuvant Chemotherapy Regimens. *Oncology Nursing Forum*, 36 (5):563-570
- Berndt E, Kallich, J, McDermott A, Xu X, Lee H, Glaspy J (2005). Reductions in Anemia and Cancer-related fatigue are Associated with Improvements in Productivity in Cancer Patients Receiving Chemotherapy. *Pharmacoeconomics*; 23(5):505-514
- Blanney J, Lowe-Strong A, Rankin J, Campbell A, Allen J, Cracey J (2010). The Cancer Rehabilitation Journey: Barriers to and Facilitators of Exercise Among Patients With Cancer-Related Fatigue. *Physical therapy*, 90(8):1135-1147
- Carey M, Burish TG (1988). Etiology and Treatment of the Psychological Side Effects Associated with Cancer Chemotherapy: A critical review and discussion. *Psychological Bulletin*, 104(3):307-325.
- Cleeland CS (2007). Symptom Burden: Multiple Symptoms and Their Impact as Patient-Reported Outcome. *J. Nat. Canc. Institute Monographs*, 37:16-21
- Dickinson GE, Clark D, Sque M (2008). Palliative Care and End of Life Issues in UK Pre-registration, undergraduate nursing programs. *Nurse Education Today*, 28:163–170
- Erickson JM, Beck S, Christian B, Dudley W, Hollen P, Albritton K. (2010). Patterns of Cancer related Fatigue in Adolescents Receiving Chemotherapy. *Oncology Nursing Forum*, 37(4): 444-455.
- Gibson F, Mulhall AB, Richardson A, Edwards JL, Ream E, Sepion BJ (2005). A Phenomenologic Study of Cancer-related fatigue in Adolescents Receiving Treatment for *Cancer. Oncology Nursing Forum*, 32:651–660.
- Hofman M, Ryan J, Figueroa-Moseley C, Jean-Pierre P, Morrow G (2007). Cancer-Related Fatigue: The Scale of the Problem. *The Oncologist*, 12:4-10.
- Hwang S, Chang V, Rue M, Kasimis B (2003). Multidimensional Independent Predictors of Cancer-Related Fatigue. *J. Pain Manag.* 26(1):604-614.
- Kuchinski AM, Reading M, Lash AA (2009). Treatment-Related Cancer-related fatigue and Exercise in Patients with Cancer: A Systematic Review. *J. Med. Surg. Nurs.*, 18(3):174-180
- Lee YH, Tsai YF, Lai YH, Tsai CM (2008). Cancer-related fatigue Experience and Coping Strategies in Taiwanese Lung Cancer Patients Receiving Chemotherapy. *J. Clin. Nurs.*, 17: 876-883
- Lucia A, Earnest C, Perez M (2003). Cancer-Related Fatigue: Can Exercise Physiology Assist Oncologists? *The Lancet Oncology*, 4:616-625.
- Miaskowski C, Dodd M, Lee K (2004). Symptom clusters: The New Frontier in Symptom Management Research. *J. Nat. Canc. Institute Monographs*, 32:17–21.
- Mustian KM, Morrow GR, Carroll JK (2007). Integrative Nonpharmacologic Behavioral Interventions for the Management of Cancer Related Fatigue. *The oncologist*; 12:52-67
- Nail LM, King KB (1987). Cancer-related fatigue A side Effect of Cancer Treatments. *Semin Oncol.*, 3:257–262.

- National Comprehensive Cancer Network (NCCN) (2012). NCCN clinical practice guide-lines in oncology: cancer-related cancer-related fatigueVI; Available at: http://www.nccn.org/professionals/physician_gls/PDF/fatigue.pdf/. Accessed april 12, 2012.
- Patric DL, Feketich SL, Frame PS, Harris JJ, Hendricks CB, Levin B. (2003). National Institutes of Health State of the Science Conference Statement: Symptom Management in Cancer: Pain, Depression, and Fatigue. *J. Nat. Canc..Institute*, 95(15):1110-1117.
- Prue G, Rankin J, Allen J, Gracey J, Cramp F. (2006). Cancer-Related Fatigue: A critical Appraisal *Eur. J. Canc.* 42(7):846-863
- Rayan JL, Carroll JK, Rayan EP, Mustian KM, Fascella K, Morrow GR (2007). Mechanisms of Cancer-Related Fatigue. *Oncol. J.* 12(1):22
- Simon A, Zittoun R (2000). Cancer-related fatigue in Cancer Patients. *Current Opinion Oncol.*; 11:244-249
- Wasteson E, Brenne E, Higginson I, Hotopf M, Ioyd-Williams M, Kaasa S.(2009). Depression Assessment and Classification in Palliative Cancer Patients: a systematic literature review. *J. Palliative Med.*, 23(8):739–75.
- Wielgus KK, Berger AM, Hertzog M (2009). Predictors of Fatigue30 Days after .Completing Anthracycline Plus Taxan Adjuvant Chemotherapy for Breast Cancer. *Oncology Nursing Forum.* 36(1):37-48
- William E, Dale G, Godley J, Neimeye R (2003). Roles for Psychologists in End-of-Life Care: Emerging Models of Practice. *American Psychological Association*, 34(6):626 – 633.
- Yeh CH, Chiang YC, Lin L, Yang CP, Chien LC, Weaver MA, Chuang HL (2008). Clinical factors associated with cancer-related fatigue over time in pediatric oncology patients receiving chemotherapy. *Brit. J. Canc.*, 99:23-2
- Yeşilbalkan OU (2009).The Effectiveness of Nursing Educationas an Intervention to Decrease Fatigue in Turkish Patients Receiving Chemotherapy.*Oncology Nursing Forum.* 36(4): E215-E222.
- Youngmee K, Jane T, Hickok MDMPH, Gary Morrow MS (2006). Cancer-related fatigue and Depression in Cancer Patients Undergoing Chemotherapy: An Emotion Approach, *J. Pain and Symptom Manag.*; 32.
- Yurtsever S (2007).The Experience of Cancer-related fatigue in Turkish Patients Receiving .Chemo therapy. *Oncology Nursing Forum*, 34(3):721-728