



Healthier eating with blanching: Retaining nutrients while enhancing taste

Shui Luo*

School of Food and Biological Engineering, Hefei University of Technology, Hefei, China

E-mail: shuiluo@hfut.edu.cn

Food emulsions play a crucial role in the texture, taste, and stability of many of the foods we consume daily. From creamy salad dressings and smooth mayonnaise to velvety ice creams and rich chocolates, emulsions are responsible for creating the desirable sensory experiences that make these products enjoyable. In this article, we will delve into the fascinating world of food emulsions, exploring their composition, preparation, properties, and applications across various culinary delights. An emulsion is a dispersion of one immiscible liquid (the dispersed phase) within another (the continuous phase), stabilized by a third component known as an emulsifier or surfactant. In food emulsions, water and oil are the most common immiscible phases, with the emulsifier acting as a bridge between the two, allowing them to remain stably combined (Du et al., 2013).

The role of emulsifiers is emulsifiers are amphiphilic molecules, meaning they possess both hydrophilic (water-loving) and hydrophobic (oil-loving) regions. This unique structure enables them to position themselves at the interface between water and oil, reducing the surface tension and preventing the coalescence of the dispersed phase. Common food emulsifiers include lecithin, mono- and diglycerides, and polysorbates, among others. Food emulsions can be classified into two main types based on the continuous phase: oil-in-water (o/w) emulsions: in these emulsions, oil droplets are dispersed within a continuous water phase. Examples include milk, salad dressings, and vinaigrettes. Water-in-oil (w/o) emulsions: in w/o emulsions, water droplets are dispersed within a continuous oil phase. Examples include butter, margarine, and cream (Kagawa et al., 2013).

Formulation and stability of food emulsions is creating stable food emulsions is a delicate process, involving the proper selection of emulsifiers and understanding the

physicochemical properties of the ingredients. Factors such as temperature, pH, and shear forces during processing can affect emulsion stability. Understanding the principles of emulsion science is vital for food manufacturers to produce products with consistent quality and extended shelf life. Emulsification technique - various techniques are employed to create food emulsions, each suited to different applications and processing conditions. Some common methods include: high-pressure homogenization: this process involves passing the emulsion through a high-pressure homogenizer, which breaks down the droplets to smaller, more stable sizes, mechanical stirring: simple agitation using mixers or blenders can create small emulsion droplets, ultrasonication: high-frequency sound waves are used to create small droplets and enhance the stability of the emulsion, microfluidization: emulsions are processed through microchannels to achieve uniform droplet size (Liu et al., 2019).

Emulsions in culinary creations- Food emulsions are integral to a wide range of food products, contributing to their texture, mouthfeel, and appearance. Some common examples include: mayonnaise: Oil-in-water emulsions that combines oil, egg yolk, and vinegar or lemon juice to create a thick, creamy texture. Ice cream: water and air are emulsified in a continuous fat phase to give ice cream its smooth and creamy texture. Salad dressings: emulsions of oil and vinegar are stabilized with an emulsifier to prevent separation and achieve a smooth consistency, chocolate: the cocoa butter in chocolate acts as an emulsifier, ensuring a smooth texture and preventing cocoa solids from settling (Peng et al., 2018).

Challenges in food emulsion technology are while food emulsions have transformed the culinary world, they present some challenges in terms of stability and sensory attributes. Issues like creaming (where droplets rise or settle), coalescence (droplet fusion), and phase separation

Received: 22-July-2023, Manuscript No. AJFST-23-110846; **Editor assigned:** 26-July-2023, Pre QC No. AJFST-110846 (PQ); **Reviewed:** 08-Aug-2023, QC No. AJFST-23-110846; **Revised:** 17-Aug-2023, Manuscript No. AJFST-23-110846 (R); **Published:** 22-Aug-2023

Citation: Luo (2023). Healthier eating with blanching: Retaining nutrients while enhancing taste. AJFST: 038.

(e.g., oil and water separation) can impact product quality. Food scientists continually strive to develop innovative emulsification techniques and emulsifiers to overcome these challenges and improve product stability.

Food emulsions are at the heart of countless culinary delights, enriching our dining experiences with their smooth textures and delightful tastes. Understanding the science behind emulsions allows food manufacturers to create products that captivate our senses and deliver consistent quality. As research in food emulsion technology continues to progress, we can expect even more delectable innovations that redefine the way we enjoy our favourite foods (Vel et al., 2019).

References

- Du Y, Shi S, Jiang Y, Xiong H, Woo MW, et al (2013). Physicochemical properties and emulsion stabilization of rice dreg glutelin conjugated with kcarrageenan through Maillard reaction. *J Sci Food Agric.* 93: 125-33.
- Kagawa Y, Maeda T, Kato Y, Ueda I, Kudo T, et al (2013). Influence of the slow infusion of a soybean oil emulsion on plasma cytokines and ex vivo T cell proliferation after an esophagectomy. *J Parenter Enteral Nutr.* 37: 123-8.
- Liu H, Cui SW, Chen M, Li Y, Liang R, et al (2019). Protective approaches and mechanisms of microencapsulation to the survival of probiotic bacteria during processing, storage and gastrointestinal digestion: A review. *Crit Rev Food Sci Nutrition.* 59: 2863-78.
- Peng C, Svirskis D, Lee SJ, Oey I, Kwak HS, et al (2018). Design of microemulsion system suitable for the oral delivery of poorly aqueous soluble beta-carotene. *Pharm Dev Technol.* 23: 682-8.
- Vel E, Sampers I, Raes K (2019). A review on influencing factors on the minimum inhibitory concentration of essential oils. *Crit Rev Food Sci Nutr.* 59: 357-78.