

Full Length Research Paper

Some aspect of dormancy studies and vitamin D content in four tree seed species

¹Ajiboye A.A*, and ²Agboola D.A

¹Department of Biological Sciences, Osun State University, Osogbo, Osun State, Nigeria.

²Department of Biological Sciences, University of Agriculture, Abeokuta, Ogun State, Nigeria

Accepted 7 February 2011

Selected leguminous tree seeds of *Tamarindus indica* (L) ,*Parkia biglobossa* (Jacq) R . Br . Don , *Prosopis africana* (Guill) Perr & Taub)and *Albizia lebbbeck* (Linn) Benth play prominent roles in nutrient recycling of soils in both savanna and tropics of Nigeria. These trees are recklessly felled without a corresponding afforestation programme. Therefore, the danger of being extinct is imminent. However the seeds of these trees were found to be dormant. The viability test used was floating methods. Efforts to alleviate their dormancy led to the usage of some dormancy-breaking methods which involve chemical scarification of concentrated sulphuric acid at 5min,10min and 15min.The viability test of the seeds showed 90-100%.Initial germination trials gave 0-20% in all the seeds.About 80-100% germination was observed when treated with concentrated sulphuric acid under 5mins duration within 10days.At 10min duration the treatment was 100% in *A.lebbbeck* while maximum percentage germination were obtained in *A. lebbbeck* (100%) and *P. biglobossa* (60%) at 15min duration. The highest vitamin D content was found in *P.africana*. This work will assist in the propagation of these valuable leguminous seeds; and reveal the potentials of these tree seeds for commercial and domestic purposes.

Key words: Dormancy, Vitamin.D, Germination, *Prosopis africana*, *Tamarindus indica*, *Parkia biglobossa*, *Albizia lebbbeck*

INTRODUCTION

Dormancy in seed is regarded as a state whereby an intact and viable seed fails to germinate or to complete germination under favorable conditions. The seeds of some leguminous plants are prevented from completing germination because the embryos are sometime constrained by its surrounding structures. This phenomenon could be termed coat-enhanced dormancy state (Ajiboye , 2009). Embryos isolated from these plants are not mostly dormant. A second category of dormancy is found in which the embryo itself is dormant, in such that it does not support the completion of germination. This state could be termed embryo-enhanced – dormancy (Ajiboye ,2009).

The Studying of germination may be difficult because population of seeds do not complete the process. Synchronous release from dormancy may be more erratic because of the threshold stimulus require to promote

germination varies widely among individual seeds. A “biotime” concept has been introduced which incorporates mathematical model to characterize and predict seed germination behaviour with respect to dormancy and the factors that influence it (Brandford,1996). Is dormancy the result of a deficiency in some vital cellular event of germination? Is there some dormancy imposed event that must be negated before germination can be completed? A broader issue is weather release from dormancy could be triggered by a variety of environmental and chemical stimulus is mediated through a common signal transduction chain that coordinates diverse cellular responses but may differ between the seeds of different species and dormancy types (Vleeshouwers *et al*, 1995). It has been suggested that there are related or common receptors for dormancy breaking agents within the plasma of the responsive embryonic cells. When triggered, these receptors then initiate a signal transduction cascade, perhaps involving synthesis of or sensitization to germination-promoting

*Corresponding author Email: abidupsy@yahoo.com

Table 1. Percentage Viability Test

Tree Seed Species	Percentage Viability (%)
<i>Tamarindus indica</i>	100
<i>Parkia biglobossa</i>	90
<i>Albizia lebbek</i>	100
<i>Prosopis Africana</i>	100

giberrellins (GAs) that leads to the completion of germination (Hilhorst, 1993; Vleeshouvers *et al*, 1995). Changes in the phosphorylating activity of membrane associated Ca^{2+} dependent protein kinases that lead to dormancy or germination has been proposed as well (Trawavas, 1988).

Vitamins are organic compounds required in trace amount in the diet for healthy growth and reproduction. They are natural materials that can be isolated from biological organisms (Bieri, 1976). The vitamin D are classified as one of the fat-soluble vitamins, because of their ability to dissolve in organic solvents. It can be transported and stored for long period of time in a manner generally similar to that of fats. Unlike the water-soluble vitamins, the fat-soluble vitamins generally do not function as co-enzymes and rarely utilized by microorganisms (Carpenter, 1974). Vitamin helps to increase the calcium absorption and assist in bone growth and teeth development.

This work focus on the dormancy studies and vitamin D contents of valuable leguminous tree seed species in Nigeria.

MATERIAL AND METHODS

Seed collection and Processing

Matured seeds were collected and extracted from their parent plant after fruit fall. Pods of *Tamarindus* and *Parkia* seeds were soaked for 24 hours while that of *Prosopis* and *Albizia* were hit with gentle strokes of moderate size stone. Dried pods of the latter were collected from the campus of the University of Ilorin, Ilorin, Kwara State of Nigeria. The seeds of *Tamarindus* were bought in Murbi market of Adamawa State, Nigeria. The dried pulps of *Parkia* were collected from its tree found at the back of female hostel of the University of Agriculture, Abeokuta, Ogun state, Nigeria.

Seed viability study

The viability of the seeds was carried using simple floating test. Seeds were immersed in distilled water contained in 500ml beaker. Those seeds found floated were considered to be damaged or bad for germination. Such seeds were discarded. Whereas, those that settled at the bottom of the water were termed to be viable for germination. Such seeds were put in container and placed in the refrigerator set at 15-20°C. However, the percentage germination was observed at each case.

Initial germination studies

About ten seeds were randomly selected at each case, after being surface sterilized with 0.1% of mercuric chloride ($HgCl_2$) for 30secs and rinsed in several changes of distilled water. The seeds were slated for germination in 9cm in diameter Petridishes containing two sterile filter papers soaked with distilled water. Three replicates of the set-up were made in this and subsequent experiment. The experiment was monitored and observed for germination for a period of 10-20days in the Botany/Zoology laboratory.

Chemical scarification

Seeds were prepared in a lot at each case with respect to species used, and immersed in concentrated Hydrogen Tetraoxosulphate VI acid for the period ranging from 5min, 10min and 15min. The set-up was raised in several changes of distilled water and air dried at room temperature ($30 \pm 20^\circ C$).

Vitamin extraction

Approximately 0.125 g of sample was measured into a set of clean test tubes and marked to 500ug volume with distilled water. About 10g/L of ascorbic acid was added as an antioxidant and shake for 15min followed by 5mins of sonication. Triton of 0.5g/L was added as detergent and 400uL acetone nitrile was added as well and mixed thoroughly, then 400uL n-hexane was measured and added and shook vigorously for 4 units. The mixture was centrifuged for 2min at 800RPM. The supernatant for vitamins determination was collected over HPLC. Separation of vitamin was carried out using BDS hypersil CN 150min, 5µm column in combination with a Javelly NH_2 guard column at wavelength 325 (Cook, 1998).

RESULTS

Table 1 showed percentage viability of various leguminous tree seed species after "floating testing". The *T.indica*, *A.lebbeck* and *P. africana* had 100% germination. *P.biglobossa* gave 90% germination.

The poor germinability of the leguminous seeds were exhibited in the initial studies shown in Table 2. Seeds prepared for germination without prior germination treatments that is, breaking the barrier of germination had reduced the rate of germination to 0-20%, 0-10%, 0-10%, and 0-10% for *T.indica*, *P.biglobossa*, *A.lebbeck* and *P.biglobossa* respectively (Table 2).

Treatment of the seeds with concentrated sulphuric acid gave rise to the percentage germination of the seeds. This is evident in the in *Albizia*, *Parkia* and *Tamarindus*

Tree Seed Species	Percentage germination (%)
<i>Tamarindus indica</i>	0-20
<i>Parkia biglobossa</i>	0-10
<i>Albizia lebbbeck</i>	0-10
<i>Prosopis Africana</i>	0-10




Figure 1. Effect of sulphuric acid scarification on germination of leguminous tree seeds (at 5mins duration).




Figure 2. Effect of Sulphuric acid scarification on germination of leguminous tree seeds at 10mins duration.

having 80-100% germination under 5min duration treatments. The control showed 0% germination within 10days (Figure 1).

Figure 2, showed at 10min, the treatment of concentrated sulphuric acid and its treatment on percentage germination of the leguminous seeds. *A.lebbbeck* seeds showed percentage germination of 100%. However, 20% germination was showed for *P. africana* and *P.biglobossa*

at each case while *T. indica* and control were 0% germination within 10days.

At 5min scarification duration, maximum percentage germination of the seeds were observed to be 100% and 60% in *A. lebbbeck* and *P. biglobossa* respectively. However, *P.africana*, *T.indica* and control showed 0% germination within 10days (Figure 3).

The highest vitamin D content of the seeds was found in




Figure 3. Effect of sulphuric acid scarification on germination of leguminous tree seeds at 15mins duration.




Figure 4. Vitamin D content of Four Leguminous Tree Seed species

P. africana having 0.8 μ g/ml while *T.indica* , *P.biglobossa* and *A. lebbeck* gave 0.7 μ g/ml, 0.75 μ g/ml and 0.75 μ g/ml respectively (Figure 4).

DISCUSSION

Viability remained high in *T.indica* , *P.biglobossa* , *P.africana* and *A.lebbeck* seeds. This implied that the embryos are in good state and that the physiological being of these seeds were not affected. Therefore, the seeds were in readiness for germination to take place, provided all the conditions necessary for germination were present.

The initial germination test revealed that the seeds exhibited physical dormancy due to hardness of the seed coats (Ajiboye , 2009) , when raised for germination without pre-germination treatments. This had led to high reduction of their germination potentials. Seeds may exhibit one form of dormancy or the other which has to do with the inability of water and other important materials

such as gases and other metabolites to pass into the embryo through the seed coats . The seed coats of dormant seeds prevented the influx of these materials to mobilize the food reserves in the embryo via imbibition of water in it that may lead radicle protrusion and later plumule(Ajiboye,2006). These are evidence of germination in viable and dormant-free seed. Some seeds lose their dormancy while in dry state when the rate of metabolism is very low. However, imbibed dormant seeds are metabolically active and this may receive an external signal like light, chilling alternating temperature and chemical or hormonal treatments that can stimulate germination (Bewley, 1997).

Termination of dormancy in seeds of *T.indica*, *P.biglobossa*, *A.lebbeck* and *P.africana* occurred by pre-treating them with concentrated sulphuric acid as a pre-sowing agent to rupture the surrounding walls of the hard seed coats after 5min,10min and 15 min of duration. The primary events in the release from dormancy are the reception of stimulus by the reception by the embryo and the immediate signal transduction chain that led to the

secondary events which could involve metabolic and hormonal changes (Bewley, 1997). The result is emergence of the embryo axis from the seeds, that is the completion of germination. (Bewley, 1997). *P. africana* showed the highest content of vitamin D (0.8ug/ml) content. This implied that the seeds are relatively rich in its content. Therefore it could be used as alternative for vitamin D in daily diets. Vitamin D is good in improving healthy growth and development. The seeds could therefore be used as a substitute for vitamin D in daily diets.

REFERENCES

- Ajiboye AA (2009). Seed germination and Seedling Physiology of Four Multipurpose Savanna Tree Species; P.hD Thesis, University of Agriculture, Abeokuta, Ogun State, Nigeria. 351pp.
- Ajiboye A.A and Ebofin, A.O; Atayese; M A; Adedire; M.O and D.A Agboola (2006): Effect of Presowing Treatment and Ectomycorrhiza on Seed Germination and Seedling Growth of Two Multipurpose Savanna Tree Legumes, *Dialium guineensis* and *Prosopis africana*. Biol. Environ. Sci. J. Tropics (BEST) 3(4):23-28.
- Bewley JA (1997): Seed germination and Dormancy, *Plant cell* 9:1055-1066
- Bieri JG, Farrell PM (1976). Vitamin E in: Vitamin and Hormones, vol 34, P.L Muson, J Glover, Ex Diczfalvsi et al, Eds. New York Academic Press pp 31.
- Cook JD, Fowers CH, Skikne BS (1998): An assessment of dried blood-spot technology for identifying iron deficiency Blood . 92: 1807-1813
- Carpenter, M.P, Howard, C.N, Jr (1974): Vitamin steroids and liver microsomal hydroxylations; Ann. J. Clin. Nutri. 27:966.
- Hilhorst HWM (1993): New aspects of seed dormancy in fourth international workshop on seeds, Basic Applied Aspects of Seed Biology D.com and F.Corbineau, eds .Pp571-579.
- Vleeshouwers LM, Bonwmeester HJ, Karsaen ?????(1995): An attempt to integrate Physiology and Ecology ; J. Ecol. 83:1031-1037