The Relationship between Quality of Life and Supportive Care Needs for Gynaecological Cancer Patients Undergoing Chemotherapy in a Regional Teaching Hospital at Southern Taiwan

Sheng-Hui Su

Supervisor, Department of Nursing, Yuan's General Hospital

Abstracts

Purpose To evaluate the relationship between quality of life and supportive care needs and investigate the most frequent unmet supportive care needs in women with gynaecological cancer during chemotherapy in a regional teaching hospital.

Method This longitudinal study was conducted on 18 patients underwent a 6-course chemotherapy from March 2018 to August 2018. The information of Quality of Life (EO-RTC-QLQ-OV28) and the Support Care Needs Survey Summary (SCNS-SF34) were obtained after each course of chemotherapy. Totally 6 sets of questionnaires were collected during the whole course of study period. The correlation of data obtained from quality of life versus those from SCNS-SF34 was studied using linear regression analysis.

Result/This study found that there was a significant negative relationship between the support care needs and the quality of life (r = .0.777, p < 0.01). The top three unmet needs of the supportive care were attitude to disease /treatment, other chemotherapy side effects, and body image.

Conclusion This study demonstrated that during the course of chemotherapy, the extent of supportive care needs varied significantly depending on the patient's requirements. The unmet supportive care needs should be identified early at the time of chemotherapy. We should provide individual care depending upon patients' needs, which may contribute to the improvement of the quality of care in patients with gynaecological cancer.

Biography:

Sheng-Hui Su has completed her BSN at Fuying University and Studies for her master at the department of Nursing Fuying University, Taiwan. She is the nursing supervisor at



Yuan's General Hospital, a regional teaching hospital. She has been involved in clinical nursing research.

Recent Publications:

- Guenther, J., Stiles, A., & Champion, J. D. (2012). The lived experience of ovarian cancer: a phenomenological approach. Journal of American Academy Nurse Practitioners, 24(10), 595-603. doi: 10.1111/j.1745-7599.2012.00732.x
- 2. Goker, A., Guvenal, T., Yanikkerem, E., Turhan, A., & Koyuncu, F. M. (2011) Quality of life in women with gynecologic cancer in Turkey. Asian Pacific Journal of Cancer Prevention, 12(11), 3121-3128.
- 3. Reese, J. B., Blackford, A., Sussman, J., Okuyama, T., Akechi, T., Bainbridge, D., & Snyder, C. F. (2015). Cancer patients' function, symptoms and supportive care needs: a latent class analysis across cultures. Quality of Life Research, 24(1), 135-146.
- Stavraka C,Ford A, Ghaem-Maghami S, et al. A study of Symptoms described by ovarian cancer survivors. Gynecol Oncology. (2012);125:59Y64
- 5. Sheng-Hui Su et al; Public Health Nursing; 2012 Feb 5.

5th Nursing Research and Evidence Based Practice Conference

Citation: Sheng-Hui Su; The Relationship between Quality of Life and Supportive Care Needs for Gynaecological Cancer Patients Undergoing Chemotherapy in a Regional Teaching Hospital at Southern Taiwan; Nursing Research 2020; April 17, 2020; Singapore City, Singapore.

J. Res. Nurs. Midwifery 2020 Volume and Issue: S(1)